

TOPOGRAPHIC GENOTYPING

FIELD OF THE INVENTION

The present invention relates to the detection of diseases, including cancers or heredity based defects in patients.

5 More specifically, the present invention relates to detecting diseases in patients from a biological specimen based on histopathologic characteristics of the specimens as observed on the microscopic level.

BACKGROUND OF THE INVENTION

10 Recent advances in genetic research, especially those focused upon cancer or inherited disorders, has led to the identification of new genes having specific patterns of DNA sequence alterations directly related to pathologic disease states (Weinberg RA, Oncogenes, Antioncogenes, and the Molecular Basis of
15 Multistep Carcinogenesis. Cancer Res., 49:3713-3721, 1989). Central to this research is the Human Genome Project, a monumental world-wide scientific effort to fully map and sequence the human genome (Watson JD, The Human Genome Project: Past, Present, and Future. Science, 248:44-49, 1990). Together the results will
20 provide a detailed blueprint of the normal Human genome together with a understanding of DNA damage upon which the diagnosis and treatment of many conditions may be formulated. Using this information, genetic based therapies have already been instituted, consisting of the introduction into selected cells of normal or
25 modified human genes designed to integrate and function as part of the host genome (Anderson WF, Human Gene Therapy. Science, 256:808-813, 1992). These initiatives provide a strong stimulus for tissue based methods which can characterize in detail DNA sequence alterations in selected cellular components of normal and disease
30 affected human tissues.

To realize the potential of the expanding database of DNA sequence information, it has become necessary to have available methods which can detect and characterize DNA sequence alterations in tissue specimens such as those routinely obtained during the medical management of patients. Presently, in clinical practice, genetic analysis usually requires a fresh and relatively large tissue sample secured independently of other specimens for diagnostic purposes. Realistically, many clinical specimens, for which genetic sequence information would be vitally needed, are of small size obtained through biopsy procedures. Moreover there exists a priority of tissue management in that proper histopathologic diagnosis is paramount demanding that adequate tissue first be secured and placed into appropriate fixative solutions to preserve morphologic integrity for accurate histopathologic evaluation. Standard practices of genetic analysis are generally ineffective on specimens exposed to fixative agents (Ben Ezra J, Johnson DA, Rossi J, Cook N, Wu A, Effect of Fixation on the Amplification of Nucleic acids from Paraffin-Embedded Material by the Polymerase Chain Reaction. J. Histochem. Cytochem., 39:351-354, 1991). This has led some to the collection of fresh tissue in Freezer Banks, a process that is inconvenient, costly and very often not carried out in practice. In contrast stands the vast bulk of archival tissue specimens in the form of fixative treated, paraffin embedded tissue blocks. These tissue archives are present in all medical centers and contain valuable specimens from patients whose disease has been followed for many years as part of the normal clinical management. These informative specimens await a simple, effective means for their genetic analysis. Clinical practice is very much in need of easily applicable techniques for DNA sequence analysis from routinely prepared tissue blocks (Antonarakis SE, Diagnosis of Genetic Disorders at the DNA Level. N. Engl. J. Med., 320:153-163, 1989). The methods to be used should not be destructive to the blocks and

must take advantage of the important insights gained through detailed histopathologic analysis. The techniques should be independent of archival storage time permitting the opportunity for long-term retrospective study. Finally the approach should be cost effective and timely to participate in everyday clinical decision making.

Topographic Genotyping (TG) is a novel system of tissue management comprehensive in scope specifically developed to address these specific issues enabling full DNA analysis within the context of traditional pathology. TG permits tissue specimens, routinely fixed in standard fixative chemical agents, of any size including minute needle biopsy specimens and cell blocks of cytology material, and of any age including those stored in paraffin for over thirty years, to be both fully available for standard histopathology examination as well as DNA sequence analysis. Furthermore, TG has been specifically designed to incorporate procedures for tissue and information handling allowing quick and easy clinical as well as research application. In essence TG is designed to allow the user to simply and effectively sample minute morphologic targets within fixative treated tissue specimens based on histopathologic and topographic considerations, which in turn may serve as the basis for detailed DNA sequence analysis. The results of TG is an integration of genetic and histopathologic features in a simple, reliable and cost effective manner for clinical application.

Solid tissue specimens, removed at surgery or through biopsy procedures, are exposed to fixative agents designed to prevent tissue breakdown and preserve morphologic integrity for microscopic analysis and archival storage. Fixatives, the most common being a 4% buffered solution of formaldehyde, cause their tissue preserving effect by a process of chemical crosslinking of

cellular constituents including proteins, sugars and nucleic acids. Much of the tissue stabilizing effect of tissue fixatives is chemically irreversible (Greer CE, Oeterson SL, Kiviat NB, Manos MM, PCR Amplification from Paraffin-Embedded Tissues. Effect of
5 Fixative and Fixation Time. Am. J. Clin. Path., 95:117-124, 1991). This tissue stabilizing and preserving chemical interaction, essential for microscopic analysis, greatly interferes with the manipulation of DNA for genetic investigation representing a major deterrent for general application of molecular analysis on fixed
10 tissue specimens.

In order to meet the need for up to date genetic analysis, current medical practice recommends obtaining separate tissue specimens, not subject to chemical fixation, exclusively for the purpose of genetic analysis. When this involves a fluid specimen of homogeneous character such as a blood sample or bone marrow aspirate, division of the specimen for separate microscopic and molecular biologic analysis is usually accomplished fairly easily without involving interfering with traditional pathologic diagnosis. For many needle biopsy procedures, however, and in a variety of other circumstances of limiting tissue availability, a solid tissue sample will not be able to be appropriately divided and thus molecular examination would not be performed. Even in the case of large specimens, which might appear at first to provide generous amounts of tissue sample for genetic study, appropriate
25 subdivision may not be feasible in as much as cellular heterogeneity cannot be fully appreciated until full tissue fixation and histopathologic examination is first performed. To derive the greatest benefit from genetic analysis, it is highly desirable to focus molecular analysis on selected tissue targets
30 reflecting the cellular basis of disease processes. This in turn can only be achieved following thorough histopathologic examination. This is the essential condition that must be met if

true and effective integration of pathology and molecular biology is to be achieved.

These realities provide a strong impetus to define new ways in which fixative treated tissue specimens should be handled to allow DNA structure analysis. Current protocols in this regard, while they may be available, are, in general, highly inefficient, difficult to apply widely in clinical practice and do not take histological considerations fully into account (Shibata DK, Arnheim N, Martin WJ, Detection of Human Papilloma Virus in Paraffin-Embedded Tissue Using the Polymerase Chain Reaction. J. Exp. Med., 167:225-230, 1988; Wright DK, Manos MM, Sample Preparation from Paraffin-Embedded Tissues. in PCR Protocols: A Guide to Methods and Applications. Innes MA, Gelfand DH, Sninsky JJ, White TJ (eds). pp. 153-158, 1990, Academic Press, Berkeley, CA; Greer CE, Lund JK, Manos MM, PCR Amplification from Paraffin-Embedded Tissues: Recommendations on the Fixatives for Long-Term Storage and Prospective Studies. PCR Meth. & Applic. 1:46-50, 1991). There exists at present no systematic means to integrated morphologic and genetic analysis of solid tissue specimens. Authoritative sources have recommended a system of DNA extraction and precipitation analogous to that used with fresh tissue (Shibata DK, Arnheim N, Martin WJ, Detection of Human Papilloma Virus in Paraffin-Embedded Tissue Using the Polymerase Chain Reaction. J. Exp. Med., 167:225-230, 1988). Nucleic acid precipitation from fixative treated specimens is very inefficient with resultant low recovery yields for subsequent genetic analysis (Ben Ezra J, Johnson DA, Rossi J, Cook N, Wu A, Effect of Fixation on the Amplification of Nucleic acids from Paraffin-Embedded Material by the Polymerase Chain Reaction. J. Histochem. Cytochem., 39:351-354, 1991). This in turn demands sacrifice of large amounts of starting material which is impractical and highly undesirable in many instances. Genetic analysis of fixative treated tissues often require gross

dissection of the paraffin block in the dry state, a procedure that is both uncontrolled, wasteful and destructive to the archival stored tissue. This approach fails to take into account important microscopic features since handling of the tissue in the paraffin block with the naked eye is not subject to fine microscopic control. Protocols direct one to scrape tissue off the slide with a scalpel. Occasionally protocols suggest selecting the area according to structures seen using the human eye (not microscope) similarly ignore the benefits derived from careful histopathologic/topographic selection (Greer CE, Lund JK, Manos MM, PCR Amplification from Paraffin-Embedded Tissues: Recommendations on the Fixatives for Long-Term Storage and Prospective Studies. PCR Meth. & Applic. 1:46-50, 1991). Tissue sampling must be performed at the microscopic level responding to the unique topographic and histopathologic features that are present in an individual tissue specimen.

DNA extracted from fixative treated tissues is generally regarded as a relatively poor starting material for nucleic acid amplification, mutational analysis and DNA sequencing (Ben Ezra J, Johnson DA, Rossi J, Cook N, Wu A, Effect of Fixation on the Amplification of Nucleic acids from Paraffin-Embedded Material by the Polymerase Chain Reaction. J. Histochem. Cytochem., 39:351-354, 1991). Users are cautioned to this effect and advised to expect poor results. Yet in selected circumstances, these very same specimens may be shown to yield important genetic information indicating that given the right approach such tissue may be very informative. Despite the knowledge that specific DNA sequence alterations are frequently found involving oncogene and tumor suppressor genes commonly present in many forms of human cancer, there exists at present no effective means to broadly analyze fixative treated specimens of any size and age for clinical application of genetic information.

Topographic genotyping was developed to meet the needs for selection of fixed tissue for genetic analysis. The specific criteria upon which topographic genotyping is based are outlined in Table 1. TG is the only system at present which fully meets these
5 necessary criteria for clinical application. TG furthermore includes the necessary organization of methodological steps and information flow for suitable clinical application at this time. While reports by others document the use of fixative treated tissues for genetic analysis none is uniquely designed to be fully
10 integrated into traditional histopathology in a simple, reliable, efficient and cost effective fashion. Criteria for the topographic component as outline in Table 1 provide the essential link to merge modern genetic analysis into traditional pathology practice.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2

TABLE 1

TOPOGRAPHIC GENOTYPING: TOPOGRAPHIC COMPONENT
CRITERIA FOR SUCCESSFUL APPLICATION

BASIC CRITERIA

EXPANDED CRITERIA

5

EFFECTIVE ON FORMALIN-FIXED, PARAFFIN-
EMBEDDED TISSUES WITHOUT UNDUE
SPECIMENS SACRIFICE

EFFECTIVE WITH DIFFERENT TYPES OF
FIXATIVES

ARCHIVAL TISSUE BLOCKS OF VARYING
STORAGE AGE

HISTOPATHOLOGIC &
TOPOGRAPHIC SAMPLING GUIDED BY
MICROSCOPIC CONSIDERATIONS

DETAILED SEPARATION OF MIXTURES OF
CELLULAR COMPONENTS

SAMPLING OF NECROTIC, CRUSHED OR
OTHERWISE UNSUITABLE SITES

CONFINED TARGETING INVOLVING MINUTE
BUT HIGHLY REPRESENTATIVE TISSUE
SAMPLING

BIOPSY SIZED SPECIMEN HANDLING

15

CAPACITY FOR MULTIPLEX GENETIC ASSAYS
WITHOUT THE NEED TO RE-CUT OR FURTHER
SACRIFICE TISSUE

COST EFFECTIVENESS OF TESTING QUALITY ASSURANCE AND CONTROL

SUMMARY OF THE INVENTION

10 The present invention pertains to a method for
topographic genotyping. The method comprises the steps of placing
a biological specimen having DNA of a patient under a microscope.
5 Then there is the step of inspecting the biological specimen
microscopically with the microscope. Next there is the step of
choosing a microscope size target on the biological specimen based
on its histopathologic characteristics. Next there is the step of
separating the target from the specimen. Then there is the step of
obtaining DNA sequences from the target so the DNA sequences can be
amplified. Next there is the step of amplifying the DNA sequences.
Then there is the step of detecting mutations in the DNA sequences.

15 The present invention pertains to a method for
topographic genotyping. The method comprises the steps of
separating a section from a specimen of fixative treated tissue.
Then there is the step of obtaining DNA sequences from the section.
Next there is the step of amplifying the DNA sequences by cyphaling
them in a PCR machine, with each cycle heating them to a
temperature no greater than 99°C, and then back to a temperature of
20 55°C in 5 minutes. Next there is the step of detecting mutations
in the DNA sequences. Preferably, the separating step includes the
step of cutting one to three 2-6 micron thick histologic sections
from the specimen.

DESCRIPTION OF THE PREFERRED EMBODIMENT

25 The present invention pertains to a method for
topographic genotyping. The method comprises the steps of placing
a biological specimen having DNA of a patient under a microscope.
Preferably, the biological specimen includes tissue sections,
cytological fluids, filter or cellular specimens.

Then there is the step of inspecting the biological specimen microscopically with the microscope. Next there is the step of choosing a microscope size target on the biological specimen based on its histopathologic characteristics. Next there is the step of separating the target from the specimen. Preferably, the specimen is a tissue section and the separating step includes the steps of slicing the target from the section and placing the target on a glass slide. Alternatively, the specimen is a tissue section and the separating step includes the step of placing the target in a tube. Alternatively, the specimen is a filter and the separating step includes the steps of cutting an arc segment from a filter and placing the segment in a tube.

Then there is the step of obtaining DNA sequences from the target so the DNA sequences can be amplified. Preferably, the obtaining step includes the step of extracting the DNA from the target. Preferably, the extracting step includes the step of placing the target in a lysis buffer. Preferably after the step of placing the target in a lysis buffer, there is the step of adding phenol chloroform into the lysis buffer with the target. Preferably after the adding step, there is the step of separating short length fragments of DNA being less than 100 base pairs in length from the target.

Next there is the step of amplifying the DNA sequences. Preferably, the amplifying step includes the step of choosing a primer corresponding to a gene of the patient. Then there is the step of adding the primer to the DNA sequences. Next there is the step of forming polymerase chain reaction on the DNA sequences with the primer.

Then there is the step of detecting mutations in the DNA sequences. Preferably, the detecting step includes the step of

determining the DNA sequence. Preferably after the determining step, there is the step of comparing the DNA sequence with known DNA sequences corresponding to DNA regions of the target. Preferably, after the detecting step, there is the step of establishing whether the DNA sequences are associated with the cancer, and applying a treatment regime to attack the cancer. Preferably, before the applying step, there is the step of identifying a source in the patient for the cancer. Alternatively, after the detecting, there is preferably the step of establishing whether the DNA sequence is associated with a condition hazardous to the health of a patient.

The present invention pertains to a method for topographic genotyping. The method comprises the steps of separating a section from a specimen of fixative treated tissue. Then there is the step of obtaining DNA sequences from the section. Next there is the step of amplifying the DNA sequences by cycling them in a PCR machine, with each cycle heating them to a temperature no greater than 99°C, and then back to a temperature of 55°C in 5 minutes. Next there is the step of detecting mutations in the DNA sequences. Preferably, the separating step includes the step of cutting one to three 2-6 micron thick histologic sections from the specimen.

For instance, once the DNA sequences have been amplified, either the step of detecting mutations in DNA sequences can be performed, or the step of identifying the DNA sequence can be performed. In essence, the step of detecting the mutations can either be through identifying the nucleotides in the DNA sequence or through a comparative technique where the individual nucleotides of DNA sequence do not necessarily need to be identified, but simply by having a comparison which matches with a known DNA sequence which conforms to a predefined cancer, the DNA sequence

can be used to identify that the specific cancer also exists in the patient. This, of course, is applicable for not just cancer but defective genes resulting from heredity or for diseases. If it is desired to specifically identify nucleotides of the DNA sequence, then the identifications specifically can be used again to compare known DNA sequences that are associated with given cancers or diseases or genetic abnormalities to heredity. These known DNA sequences can be obtained from established libraries that maintain such information and can be searched similar to, for instance, the way fingerprints are searched, although in this case, one would not need to have a graphic match, but instead would just compare the nucleotide sequences themselves for match. Once the given abnormality associated with a specific type of cancer or disease or hereditary defect is identified, much information can be obtained, for instance, with cancers, since they have a unique DNA sequence associated with a specific type of cancer, If a breast cancer is determined in a patient, and several years later, a nodule is detected in another part of the body, then a biopsy will be taken of the nodule. The nodule will be analyzed as explained below, and its DNA sequence will be reviewed. Upon revealing or identifying its DNA sequence, it can be determined whether the source of the DNA was a cancer associated with the breast. Then the course of treatment for the nodule could be better identified since the origin of the cancer is known and it is appropriate to apply a therapy that can attack a breast cancer based origin.

Similarly, by identifying the DNA sequence and comparing it to known cancers, not only can the cancer be determined, but the stage of the cancer can be determined. If the cancer is in an early stage, then it might be indicative of chemotherapy or radiation treatment of a certain dosage or a certain chemical. If the cancer is in a later stage, it might be more indicative of a more aggressive type of therapy which might have greater side

effects but might be necessary in order to save the life of the patient. Since it is recognized that cancers do not occur essentially in periods of days or weeks but take time for the DNA sequence to mutate and the cancer to arise, it is important to
5 identify the DNA sequence to more accurately create a therapy regime which can attack and eradicate the cancer. The more information is known about the cancer or the disease, the better the therapy regime can be developed.

Topographic Genotyping (TG) refers to a multistep process
10 complementing traditional histopathology that utilizes novel adaptations allowing fixative treated specimens to be subject to detailed DNA sequence analysis (Table 2). TG extends traditional analysis by enabling the genetic analysis to be conducted on
15 microscopic sized targets selected from tissue sections based on histopathologic characteristics. Analysis can be applied to all sized specimens including archival samples. With minimal modifications, other types of clinical samples such as cytological fluids, filter and cellular specimens can be similarly analyzed. TG integrates histopathologic and genetic information in a format that is designed to be sensitive, specific, timely and cost effective
20 while at the same time not being destructive of archival tissue or competitive with established medical practices. TG is organized so that the user is constantly in touch with the progress of each test specimen which can be reported and evaluated for aspects of quality
25 assurance and control. Topographic genotyping is specifically designed to enable the microscopist to select, at leisure and in the comfort of his/her workbench, one or more microscopic sites within a tissue specimen for detailed DNA sequence analysis.

The general outline of TG is shown in Table 2. Tissues
30 undergo traditional fixative treatment as currently performed in hospital and laboratories worldwide. Microscopic sections are

prepared and stained for proper morphologic evaluation and histopathologic diagnosis. There is no interference or delay in well established pathology practices. Moreover specific aspect of histologic handling of tissues, known to vary from one institution to another such as the duration and temperature of various individual steps, exerts no detrimental effect. TG is fully compatible with routine practice variations.

One or more topographic targets are selected based on histopathologic considerations. These targets are minute in size and highly representative of the cellular alterations to which they are related. Tissue blocks are recut, baked and deparaffinized (as routinely done prior to histological staining) to allow topographic sampling directly from unstained blank sections. Minute but highly representative target samples are collected in a 0.5 mL microtube referred to as a storage unit from which up to 30 different DNA sequence analyses can be performed immediately or at any later time. This storage unit represents the link between histopathology and DNA sequence analysis enabling DNA sequence alterations to be intimately related to cellular alterations of specific disease processes. At no time are tissue blocks damaged or excessively utilized thus tissue remains available for future histological or genetic examinations.

An aliquot of the storage unit is subject to DNA amplification followed by direct sequencing to elucidate specific DNA sequence alterations. TG incorporates unique modifications optimizing DNA amplification of fixative treated DNA. Efficient and specific nucleic acid amplification is the most critical step when processing fixative treated DNA. TG has been formulated upon a scientific basis which best explains and addresses the implications of fixative treatment for DNA amplification. Greater than 95% efficiency is achieved even for old archival fixative treated,

paraffin embedded tissue blocks. DNA amplification is followed by DNA sequencing permitting comparative sequence analysis of samples within and between cellular areas within tissue specimens. Alternatively other general screening or detailed mutational analysis method may be used to search for DNA structural damage. In this way genetic information is integrated with pathologic findings. The timetable to accomplish this analysis is included in Table 2. Using the day of initial specimen receipt as 'day 1', histopathologic evaluation is generally completed on days 2-3. TG is designed to provide genetic information for integrated analysis by day 5 meeting the timely needs of clinical practice to have genetic information readily available at the earliest moment. Since TG is equally effective on biopsy specimen, genetic information can be provided based on biopsy specimens to assist planning of most involved surgical or medical procedures.

TG includes several unique aspects of special value in the clinical management of patients suspected of genetic related disease states. Tissue sampling is generally carried out only once without the need to repeatedly section the tissue block thereby avoiding unnecessary sacrifice of tissue. The topographic tissue samples can be stored indefinitely and returned to at future dates for additional genetic analyses. Appropriate elements of quality assurance and control are included for clinical laboratory institution as well as informatics organization design. TG can easily be scaled up to handle large numbers of tissue samples. TG consists of simple laboratory methods and instrumentation encouraging broad and easy application. In these ways TG represents a novel and important advance providing modern medicine with an effective tool for the handling of tissue specimens leading to better diagnosis and treatment.

TABLE 2

TOPOGRAPHIC GENOTYPING: GENERAL OUTLINE

	<u>BASIC PROCEDURE</u>	<u>ANCILLARY PROCEDURES</u>	<u>TIME- TABLE</u>
5	TISSUE FIXATION GROSS DISSECTION	PRELIMINARY SAMPLING	DAY 1
	HISTOLOGIC PREPARATION	DIRECT MINUTE TARGET SAMPLING	DAY 2
	MICROSCOPIC EVALUATION	HISTOPATHOLOGIC DIAGNOSIS TOPOGRAPHIC SITE SELECTION	DAY 2
10	TOPOGRAPHIC SAMPLING WITH CREATION OF THE TOPOGRAPHIC STORAGE UNIT	GENE SELECTION	DAY 2
	NUCLEIC ACID AMPLIFICATION	OLIGONUCLEOTIDE PRIMER DESIGN FOR DNA AMPLIFICATION AND SEQUENCING	DAY 3

GENETIC DIAGNOSIS LEADING TO DAY 4-5
INTEGRATED GENETIC/
HISTOPATHOLOGIC DIAGNOSIS

SPECIFIC DESCRIPTION OF COMPONENTS OF TOPOGRAPHIC GENOTYPING

SPECIMEN SELECTION:

Virtually any fixative-treated, paraffin-embedded tissue specimen can be effectively subject to DNA sequence analysis using TG. Standard fixative agents such as formaldehyde, Bouin's, B5, and some mercury containing solutions do not interfere with topographic genotyping. Exceptions include strong acid decalcifying solutions which cause extensive DNA damage. Undecalcified bone section may be utilized. Conditions of fixation such as duration of time between initial fixation and variations in tissue processing are not a critical factor. The user of TG is free to investigate archival fixed tissues for detailed DNA sequence alterations.

In TG, the user selects one or more microscopic sized targets irrespective of shape or size in the tissue section of interest. For example cellular alterations in very confined areas such as might be detected following the use of special procedures such as immunohistochemical staining may serve as the basis for topographic sample selection. Small, needle biopsy specimens or cytological cell blocks prove quite satisfactory. Archival age is not a detrimental factor with specimens thirty years and older being capable of analysis. Complex mixtures of tissue types including disease related and normal elements can be dissected and evaluated separately. In addition segments of cytological filter preparations or portions of cell pellets with or without fixation can be analyzed. Hence the user of TG would have the

opportunity to carry out virtually any type of retrospective analysis on specimens of his/her choice.

The microscopist would most often utilize TG when examining tissue specimens harboring or suspected to be containing microscopic evidence of cancer. In the case of large complex specimens such as surgical resections, TG would free the user from the need to obtain fresh tissue for frozen storage. Rather, the user would have the assurance that detailed DNA analysis could follow full tissue fixation. The latter is essential as a means to properly set the tissue up for detailed gross dissection which can not be effectively performed when hastily carried out on fresh unfixed tissues. Avoidance in the handling of fresh tissue is most attractive in the examination of potentially infectious tissue specimens as it lessens the risk of exposure to blood borne infections such as AIDS.

In the course of gross dissection, one or more targets can be selected based on macroscopic appearance of the carefully dissected specimen. These macroscopic selected specimens are taken in a preliminary fashion as full topographic and histopathologic significance would be confirmed by microscopic analysis. Supplemental topographic sampling may be required based on precise microscopic features. These preliminary samples could begin the genotyping analysis with results available on the fourth day by commencing the genetic component one day earlier (Table 2). This variation on the use of TG to analyze selected tissue specimens from the gross fixed specimen has broad application by encouraging consideration and incorporation of genetic evaluation into traditional pathology practice. At the end of a typical working day, the gross dissector may conveniently accumulate a series of individual samples that can then be processed in a batch for specific DNA sequence

alterations without significant time delay to standard gross fixed tissue dissection and analysis. The representativeness of the sample obtained at gross dissection would be subsequently judged at the time of microscopic evaluation at which point additional topographic targets may be chosen as required.

At the time of microscopic evaluation, the microscopist can simply place one or more ink dot marks on each site he/she wishes to analyze for DNA sequence alterations. The tissue blocks are retrieved and 1-4 four micron thick sections are cut as is general routine for typical recuts. In the case of minute discrete targets such as biopsy specimens, sampling takes place directly from the waterbath as described below. The sampled tissue is collected in 0.5 mL microtubes referred to as storage units which may be returned to for repeated genetic analyses without the need to recut the tissue block. The approach for target selection is again shown to be very broad enabling virtually any tissue specimen of any size to be subject to integrated genetic/histopathologic analysis.

GENE SELECTION AND OLIGONUCLEOTIDE PRIMER DESIGN:

With TG, any region of a gene can be amplified provided sufficient sequence information is available upon which to formulate amplifying and sequencing primers; short DNA sequences, 18 - 30 base pair long, most easily created by means of an oligonucleotide synthesizer apparatus. These primers direct the amplification and sequencing of DNA in TG. Oligonucleotide primer pairs are usually designed to amplify a genomic region approximately 200 base pairs in length, although longer lengths can be effectively amplified from fixative treated tissues. Either amplifying primer can serve as a sequencing primer, but design and use of an internal primer may in some case be

worthwhile to achieve a clean sequencing band pattern. As sequencing will be performed by means of dideoxy chain termination with ^{35}S radionucleotide incorporation, it is important to select a radionucleotide that will be incorporated as close to the 3' end of the ultimate sequencing primer, ideally within three bases and several times within the first 10 bases. [cite?]

TG is valuable as it enables the operator to freely investigate any genetic target for DNA sequence alterations from any type of fixative treated specimen. The only requirement is a minimum of sequence information upon which to design oligonucleotide primers. This genetic information is usually found in scientific manuscripts as well as in general repositories of genetic sequence information such as GenBank and EMBL and other genetic database resources readily available through network communication services. A typical application of TG would involve a potential user selecting a series of archival tissue blocks and a gene of interest for which the sequence information would be obtained. Once oligonucleotide primers have been made, the user would search for DNA sequence alterations following the guidelines and recommendations included in TG.

HISTOLOGY PREPARATION:

Once one or more targets are selected for topographic analysis, tissue blocks are routinely sectioned at 4 microns on a microtome and floated in a water bath. For most application 1 to 4 tissue sections are sufficient to create a storage unit for 20 to 30 genetic analyses. The sections are picked up on glass slides and baked for 3 hrs. in a 55° C oven. No special precautions are required other than avoidance of contact with

post amplification products. Several sections may be placed on a single glass slide. Even minor folding within the tissue section is acceptable provided the target can be adequately visualized. Less than perfect recuts that otherwise would be discarded can be source of material for topographic selection further minimizing expenditure of tissue. With a minimum of experience, the histology technician can quickly learn to anticipate the tissue needs for TG leading to highly efficient tissue usage.

When selected targets are small such as biopsy specimens, they may be taken directly from the waterbath into the storage vial without the need to scrape directly from glass slides. Direct sampling from the waterbath is completely novel resulting in a simple yet highly efficient means utilize tiny specimens. Direct sampling does result in the inclusion of paraffin in the storage unit. This, however, is not an interfering factor since high temperature treatment of the storage unit after collection of the specimen is sufficient to free the tissue of its paraffin wax surround.

With experience, the technologist responsible for direct topographic selection from the waterbath will be able to successfully sample a wide array of specimens with this approach. Tissue remaining after direct sampling can be picked up on glass slides if desired and stained to evaluate the accuracy of direct sampling ensuring a means to assess quality assurance in direct topographic sampling.

PROTOCOL: DIRECT TOPOGRAPHIC SELECTION FROM FLOATING PARAFFIN SECTIONS

PRINCIPLE OF TEST: When a specific target is small and well localized, it can be dissected directly from the water bath using fine probes bypassing sampling from glass slides.

REAGENTS: Deionized H₂O, ice and Topographic Lysis Buffer (100 mM NaCl, 25 mM EDTA pH 8.6, 10 mM Tris pH 8.3, 0.5% NP₄₀).

INSTRUMENTATION: Microtome and blade, float water bath, paint brush, fine picks, tissue papers, and 0.5 mL tubes, 4° C refrigerator and/or -20° C freezer

Microscopist reviews slides and marks area to be sampled. Matching paraffin block is cooled on ice in preparation for microtomy.

Heat deionized water in tissue float between 50 - 55° C (above melting temperature of paraffin). Cut tissue on microtome and place ribbon of tissue in water bath. Raising the temperature partially melts the paraffin and allows the tissue to float free.

Use a fine probe to lightly touch area of interest and pluck from water. To aid in accurate selection, use a second probe to hold back unwanted tissue. Place in empty 0.5 mL tube and repeat until sample collected. Add 25 µl of Topographic Lysis Buffer 1 to tube (see below).

Residual tissue can be picked up on a glass slide and stained to confirm accuracy of sampling. This direct method is not suitable for large samples as the target tissue doesn't usually separate easily from the whole, unless well encapsulated or otherwise demarcated from surrounding unwanted tissue.

To avoid contamination between samples, clear water bath of previous samples by sweeping with tissue paper. Wipe probes clean.

VARIATION: DIRECT SAMPLING OF CYTOLOGICAL FILTER SPECIMENS

Cytology fluid specimens, in the form of cytocentrifuged or vacuum suctioned filter preps can be handled in a direct fashion. One or more cytology filter arc segments, approximately 45 degrees in size, are cut and placed in empty storage tube to which lysis buffer is added. A similar approach is used when handling cytological brushes or similar materials containing cellular material. The storage unit is then processed as described below.

TOPOGRAPHIC SAMPLING FROM GLASS SLIDES:

Topographic sampling from glass slides based upon microscopic considerations is the foundation that links genetic analysis with histopathology. It is by this detailed selection process that cellular heterogeneity may serve as the basis for integrated genetic/pathologic analysis. The unique contribution of TG is the ease by which even minute tissue targets from cellular heterogeneous specimens can be processed for multiple gene sequence alterations. With practice, even complex admixtures of cellular specimens can readily be dissected into individual units for comparative genetic evaluation.

Topographic sampling from slides is effective for virtually all tissue sections irrespective of individual histology practices. Unstained slides may be prepared in another institution then sent to a central laboratory for the genetic analysis component. The technique is not harmed by storage of undeparaffinized sections which may be kept for long periods of time and then processed.

Topographic selection of tissue based on microscopic considerations is intended to optimize sample selection and purity. Cellular characteristics as well as histochemical and immunohistochemical features serves as the basis by which the configuration of target areas are planned. The minute but highly representative nature of the tissue sample allows detailed integration of morphologic and genetic findings enhancing the overall information derived from the specimen. Even small biopsy specimens can be carefully dissected into distinct regions for comparative genetic analysis. For example in the case of a biopsy of malignant melanoma of the skin, the overlying squamous epithelium can first be removed and analyzed for genetic changes that correspond to the stage of intraepithelial growth. The superficial and deep components of an invasive tumor can be discretely sampled and analyzed separately. In this way TG represents the only means to effectively analyze such samples for DNA sequence alterations in as much as the primary skin tumor is usually small and cannot be divided for storage in a frozen tumor bank. TG would provide the means to conduct DNA sequence analysis on this specific type of tissue specimens which at present is greatly lacking. TG would ensure that abundant material would still remain in the paraffin block for future investigation.

PROTOCOL: TOPOGRAPHIC DISSECTION FROM GLASS SLIDES

PRINCIPLE OF TEST: Slides, blocks and patient data/records are reviewed. The best specimen slide or slides is/are selected, the areas of interest marked and unstained tissue sections are prepared for topographic dissection under a dissecting

5 microscope. Dissection based on histopathology allows complex mixtures of tissue to be dissected and analyzed separately. DNA damage can be related to cellular alterations. Even microscopic sized targets consisting of small collections of cells can be analyzed.

10 REAGENTS: Ice, Xylene, graded EtOH (100%, 95%, 70%), Topographic Lysis Buffer 1 (100 mM NaCl, 25 mM EDTA pH 8.6, 10 mM Tris pH 8.3, 0.5% NP₄₀), Sampling Buffer (2% NP₄₀ in H₂O). Both Topographic Lysis Buffer 1 and Sampling Buffer are stored either at room temperature or in 4°C refrigerator.

15 INSTRUMENTATION: Chemical hood, timers, staining jars, slide racks, slide holders, dissecting microscope, fine scalpel blade, sterile plastic transfer pipette, 0.5 mL sterile tube, tube racks, permanent marker, pipettor and tips, tissue float bath, microtome and blades and incubator oven, pretreated glass slides.
20 (Pretreat slides by dipping into solutions to help tissue adhere to slides. Poly-L-Lysine, silane solutions or other agents may be used to assist adherence of tissue sections (only egg albumen can not be used).

25 Microscopist reviews histology slides and marks one or more target areas. An appropriate number of unstained recuts are prepared based target size. When blocks have been cooled on ice, cut 4-8 microns on microtome and float in water bath. Pick up tissue on glass slides and label with block #. Bake 3-18 hr. in 50° C oven to optimize adherence.

Slides are deparaffinized in Xylene (5 min. X 2), 100% EtOH (5 min. X 2), 95% EtOH (5 min. X 2), 70 % EtOH (5 min. X 2). then air dried. (Limited hood space; Xylene (10 min.), then 100 % EtOH (10 min.). Change reagents often).

- 5 Arrange slides in slide holder with matching stained slide. Tissues may be left in a clean environment for several days awaiting Topographic Selection. Steps 1- 3 may be done in histology lab. To avoid contamination of samples, from this point onward all tubes, pipette tips, etc. are sterile and all solutions are made with autoclaved deionized water, unless noted.

10 Fill 0.5 mL tubes with 25 μ L Topographic Lysis Buffer (100 mM NaCl, 25 mM EDTA pH 8.6, 10 mM Tris pH 8.3, 0.5 % NP₄₀) and close. Buffer lyses cell and nuclear membranes to extract DNA.

- 15 Barely dampen target region with Sampling Buffer (a drop or bubbles), blot or air dry excess. Place slide on stage of dissecting microscope and examine under dark field optics. Use scalpel tip to scrape target site into a clump, lift off glass and place in Topographic Lysis Buffer in tube. Detergent action of sampling buffer releases tissue into buffer. Add sampling
20 buffer as needed. Repeat until sufficient sample collected then close tube.

- Highly detailed topographic dissection takes advantage of the ability of minimally wetted tissue to be cleanly separated from adjacent dry tissue. Insert clean scalpel tip into Sampling
25 Buffer. The drop of Buffer adhering to tip is then precisely placed at point of sampling. Carry this region over to the next site to continue sampling.

Typically 0.25 to 0.50 cm² area of tissue is placed in Topographic Lysis Buffer filled tube. For Quality Assurance, post-sampling slides can be stained to assess accuracy of topographic selection.

- 5 Samples may be stored in a refrigerator 4° C for up to a week, or -20° C freezer for years prior to further processing.

SCIENTIFIC BASIS FOR EFFECTIVE DNA AMPLIFICATION FROM MINUTE TOPOGRAPHIC SAMPLES OF FIXATIVE TREATED TISSUE SPECIMENS

10 The rate limiting step when handling fixative treated tissue specimens for genetic analysis is effective and specific DNA amplification. Unlike pure DNA that is extracted from fresh or frozen tissues, DNA that has been exposed to chemical fixative such as formaldehyde are often unsuccessfully or only poorly amplified. While the reason for the inability to efficiently amplify fixed tissues is known to be related to chemical action of the fixative agent upon nucleic acids, the precise mechanism most directly related to poor amplification is only poorly understood. The most suitable measures to overcome this detrimental effect are therefore not fully appreciated. Instead, 15 recourse is usually taken to sacrificing large amounts of fixative treated tissue or abandoning the use of fixative treated tissues altogether.

25 It is a misconception that fixative treated tissue provides an inadequate amount of starting DNA for nucleic acid amplification. In reality, the tissue obtained from one to three 4 micron thick histologic sections of a small biopsy specimens, when handled properly, provides sufficient material for consistent and effective DNA amplification. In fact, when one attempts to add more fixative treated tissue to the amplification

reaction, the tendency is to abolish successful amplification. This result is likely due to a combination of factors. This would include absorption of the Taq polymerase enzyme on the large amount of fixative exposed DNA that is provided by a large tissue specimen. Also errant amplification is likely when overloading the reaction with tissue and is reflected in the generation of broad smear of low and high molecular weight DNA. It is vitally important to use a small rather than a large amount of tissue to initially trigger the amplification reaction. Once triggered in the first few cycles to copy a sufficient quantity of DNA from the tissue template, the remainder of the amplification reaction goes forth in the buffer solution without significant participation of the original tissue DNA.

It is also generally held that the effect of tissue fixation is to induce irreversible cross linking of DNA and associated histone proteins that will prevent proper denaturation of the DNA in response to elevation in temperature (4), incorporated by reference. This is supported by the need to commence amplification with a series of high temperature denaturation steps early in the amplification process. This effect however is minor as demonstrated by the negative effect of high temperature annealing steps which invariably abolish amplification. In fact annealing and polymerization steps should be conducted at relatively low temperatures than that used when amplifying pure DNA. It is for this reason that two step amplification programs and shortening of the time duration of cycling program plateau phases has precisely the opposite effect than that theoretically expected. By the same token introduction of denaturing agents such as glycerol or DMSO, are found to exert no effect or have a negative impact on amplification.

It can be shown that the most important effect of tissue fixation on DNA is a loss in processivity that results from chemical modification of DNA causing the enzyme polymerization to prematurely terminate or fall off the fixative treated DNA template resulting in the production of a shorter than desired amplicon fragment. This short fragment may in a subsequent annealing step attach to an inappropriate site triggering undesirable polymerization which will result in inclusion of primer annealing sites on both ends of the nonspecific fragment. In turn this will lead to discrete nonspecific amplification bands or more often a wide range of amplified DNA reflected as a broad smear on agarose electrophoresis. It is the usual practice to raise both annealing and polymerization temperature to antagonize mispriming as discussed above but given the ease with which polymerization is terminated, this raising of temperature will have precisely the opposite effect intended. The phenomenon of premature termination is a most important factor leading to inefficient amplification and therefore measures which directly address this factor will have the most beneficial effect in the situation of fixative treated DNA.

High temperature denaturation is most valuable in the first five amplification cycles to produce maximal DNA template relaxation in anticipation of primer annealing. For most tissue specimens and gene targets this precaution together with experimentation to identify optimal functioning primers proves sufficient to achieve effective amplification. When this is found still to be inadequate, treatment of the storage unit with phenol chloroform washing as described below without the need for precipitation will confer effective amplification. This positive effect of phenol chloroform washing is due to more vigorous removal of DNA binding proteins that remain despite treatment

with proteinase K. These adherent proteins may contribute to loss of processivity.

In a small minority of cases phenol-chloroform washing will not be sufficient to allow effective specific DNA amplification.

- 5 For these infrequent cases one may resort to filter separation of small molecular weight DNA to confer effective amplification. This step is designed to separate small molecular weight short fragments of DNA from the storage tissue unit which may engage in the amplification reaction or contribute to premature termination of polymerization in the early amplification cycles. Filter separation as described below is conveniently formatted so that a minimum of manipulation is required to prepare the storage unit for amplification.

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
1000
1005
1010
1015
1020
1025
1030
1035
1040
1045
1050
1055
1060
1065
1070
1075
1080
1085
1090
1095
1100
1105
1110
1115
1120
1125
1130
1135
1140
1145
1150
1155
1160
1165
1170
1175
1180
1185
1190
1195
1200
1205
1210
1215
1220
1225
1230
1235
1240
1245
1250
1255
1260
1265
1270
1275
1280
1285
1290
1295
1300
1305
1310
1315
1320
1325
1330
1335
1340
1345
1350
1355
1360
1365
1370
1375
1380
1385
1390
1395
1400
1405
1410
1415
1420
1425
1430
1435
1440
1445
1450
1455
1460
1465
1470
1475
1480
1485
1490
1495
1500
1505
1510
1515
1520
1525
1530
1535
1540
1545
1550
1555
1560
1565
1570
1575
1580
1585
1590
1595
1600
1605
1610
1615
1620
1625
1630
1635
1640
1645
1650
1655
1660
1665
1670
1675
1680
1685
1690
1695
1700
1705
1710
1715
1720
1725
1730
1735
1740
1745
1750
1755
1760
1765
1770
1775
1780
1785
1790
1795
1800
1805
1810
1815
1820
1825
1830
1835
1840
1845
1850
1855
1860
1865
1870
1875
1880
1885
1890
1895
1900
1905
1910
1915
1920
1925
1930
1935
1940
1945
1950
1955
1960
1965
1970
1975
1980
1985
1990
1995
2000
2005
2010
2015
2020
2025
2030
2035
2040
2045
2050
2055
2060
2065
2070
2075
2080
2085
2090
2095
2100
2105
2110
2115
2120
2125
2130
2135
2140
2145
2150
2155
2160
2165
2170
2175
2180
2185
2190
2195
2200
2205
2210
2215
2220
2225
2230
2235
2240
2245
2250
2255
2260
2265
2270
2275
2280
2285
2290
2295
2300
2305
2310
2315
2320
2325
2330
2335
2340
2345
2350
2355
2360
2365
2370
2375
2380
2385
2390
2395
2400
2405
2410
2415
2420
2425
2430
2435
2440
2445
2450
2455
2460
2465
2470
2475
2480
2485
2490
2495
2500
2505
2510
2515
2520
2525
2530
2535
2540
2545
2550
2555
2560
2565
2570
2575
2580
2585
2590
2595
2600
2605
2610
2615
2620
2625
2630
2635
2640
2645
2650
2655
2660
2665
2670
2675
2680
2685
2690
2695
2700
2705
2710
2715
2720
2725
2730
2735
2740
2745
2750
2755
2760
2765
2770
2775
2780
2785
2790
2795
2800
2805
2810
2815
2820
2825
2830
2835
2840
2845
2850
2855
2860
2865
2870
2875
2880
2885
2890
2895
2900
2905
2910
2915
2920
2925
2930
2935
2940
2945
2950
2955
2960
2965
2970
2975
2980
2985
2990
2995
3000
3005
3010
3015
3020
3025
3030
3035
3040
3045
3050
3055
3060
3065
3070
3075
3080
3085
3090
3095
3100
3105
3110
3115
3120
3125
3130
3135
3140
3145
3150
3155
3160
3165
3170
3175
3180
3185
3190
3195
3200
3205
3210
3215
3220
3225
3230
3235
3240
3245
3250
3255
3260
3265
3270
3275
3280
3285
3290
3295
3300
3305
3310
3315
3320
3325
3330
3335
3340
3345
3350
3355
3360
3365
3370
3375
3380
3385
3390
3395
3400
3405
3410
3415
3420
3425
3430
3435
3440
3445
3450
3455
3460
3465
3470
3475
3480
3485
3490
3495
3500
3505
3510
3515
3520
3525
3530
3535
3540
3545
3550
3555
3560
3565
3570
3575
3580
3585
3590
3595
3600
3605
3610
3615
3620
3625
3630
3635
3640
3645
3650
3655
3660
3665
3670
3675
3680
3685
3690
3695
3700
3705
3710
3715
3720
3725
3730
3735
3740
3745
3750
3755
3760
3765
3770
3775
3780
3785
3790
3795
3800
3805
3810
3815
3820
3825
3830
3835
3840
3845
3850
3855
3860
3865
3870
3875
3880
3885
3890
3895
3900
3905
3910
3915
3920
3925
3930
3935
3940
3945
3950
3955
3960
3965
3970
3975
3980
3985
3990
3995
4000
4005
4010
4015
4020
4025
4030
4035
4040
4045
4050
4055
4060
4065
4070
4075
4080
4085
4090
4095
4100
4105
4110
4115
4120
4125
4130
4135
4140
4145
4150
4155
4160
4165
4170
4175
4180
4185
4190
4195
4200
4205
4210
4215
4220
4225
4230
4235
4240
4245
4250
4255
4260
4265
4270
4275
4280
4285
4290
4295
4300
4305
4310
4315
4320
4325
4330
4335
4340
4345
4350
4355
4360
4365
4370
4375
4380
4385
4390
4395
4400
4405
4410
4415
4420
4425
4430
4435
4440
4445
4450
4455
4460
4465
4470
4475
4480
4485
4490
4495
4500
4505
4510
4515
4520
4525
4530
4535
4540
4545
4550
4555
4560
4565
4570
4575
4580
4585
4590
4595
4600
4605
4610
4615
4620
4625
4630
4635
4640
4645
4650
4655
4660
4665
4670
4675
4680
4685
4690
4695
4700
4705
4710
4715
4720
4725
4730
4735
4740
4745
4750
4755
4760
4765
4770
4775
4780
4785
4790
4795
4800
4805
4810
4815
4820
4825
4830
4835
4840
4845
4850
4855
4860
4865
4870
4875
4880
4885
4890
4895
4900
4905
4910
4915
4920
4925
4930
4935
4940
4945
4950
4955
4960
4965
4970
4975
4980
4985
4990
4995
5000
5005
5010
5015
5020
5025
5030
5035
5040
5045
5050
5055
5060
5065
5070
5075
5080
5085
5090
5095
5100
5105
5110
5115
5120
5125
5130
5135
5140
5145
5150
5155
5160
5165
5170
5175
5180
5185
5190
5195
5200
5205
5210
5215
5220
5225
5230
5235
5240
5245
5250
5255
5260
5265
5270
5275
5280
5285
5290
5295
5300
5305
5310
5315
5320
5325
5330
5335
5340
5345
5350
5355
5360
5365
5370
5375
5380
5385
5390
5395
5400
5405
5410
5415
5420
5425
5430
5435
5440
5445
5450
5455
5460
5465
5470
5475
5480
5485
5490
5495
5500
5505
5510
5515
5520
5525
5530
5535
5540
5545
5550
5555
5560
5565
5570
5575
5580
5585
5590
5595
5600
5605
5610
5615
5620
5625
5630
5635
5640
5645
5650
5655
5660
5665
5670
5675
5680
5685
5690
5695
5700
5705
5710
5715
5720
5725
5730
5735
5740
5745
5750
5755
5760
5765
5770
5775
5780
5785
5790
5795
5800
5805
5810
5815
5820
5825
5830
5835
5840
5845
5850
5855
5860
5865
5870
5875
5880
5885
5890
5895
5900
5905
5910
5915
5920
5925
5930
5935
5940
5945
5950
5955
5960
5965
5970
5975
5980
5985
5990
5995
6000
6005
6010
6015
6020
6025
6030
6035
6040
6045
6050
6055
6060
6065
6070
6075
6080
6085
6090
6095
6100
6105
6110
6115
6120
6125
6130
6135
6140
6145
6150
6155
6160
6165
6170
6175
6180
6185
6190
6195
6200
6205
6210
6215
6220
6225
6230
6235
6240
6245
6250
6255
6260
6265
6270
6275
6280
6285
6290
6295
6300
6305
6310
6315
6320
6325
6330
6335
6340
6345
6350
6355
6360
6365
6370
6375
6380
6385
6390
6395
6400
6405
6410
6415
6420
6425
6430
6435
6440
6445
6450
6455
6460
6465
6470
6475
6480
6485
6490
6495
6500
6505
6510
6515
6520
6525
6530
6535
6540
6545
6550
6555
6560
6565
6570
6575
6580
6585
6590
6595
6600
6605
6610
6615
6620
6625
6630
6635
6640
6645
6650
6655
6660
6665
6670
6675
6680
6685
6690
6695
6700
6705
6710
6715
6720
6725
6730
6735
6740
6745
6750
6755
6760
6765
6770
6775
6780
6785
6790
6795
6800
6805
6810
6815
6820
6825
6830
6835
6840
6845
6850
6855
6860
6865
6870
6875
6880
6885
6890
6895
6900
6905
6910
6915
6920
6925
6930
6935
6940
6945
6950
6955
6960
6965
6970
6975
6980
6985
6990
6995
7000
7005
7010
7015
7020
7025
7030
7035
7040
7045
7050
7055
7060
7065
7070
7075
7080
7085
7090
7095
7100
7105
7110
7115
7120
7125
7130
7135
7140
7145
7150
7155
7160
7165
7170
7175
7180
7185
7190
7195
7200
7205
7210
7215
7220
7225
7230
7235
7240
7245
7250
7255
7260
7265
7270
7275
7280
7285
7290
7295
7300
7305
7310
7315
7320
7325
7330
7335
7340
7345
7350
7355
7360
7365
7370
7375
7380
7385
7390
7395
7400
7405
7410
7415
7420
7425
7430
7435
7440
7445
7450
7455
7460
7465
7470
7475
7480
7485
7490
7495
7500
7505
7510
7515
7520
7525
7530
7535
7540
7545
7550
7555
7560
7565
7570
7575
7580
7585
7590
7595
7600
7605
7610
7615
7620
7625
7630
7635
7640
7645
7650
7655
7660
7665
7670
7675
7680
7685
7690
7695
7700
7705
7710
7715
7720
7725
7730
7735
7740
7745
7750
7755
7760
7765
7770
7775
7780
7785
7790
7795
7800
7805
7810
7815
7820
7825
7830
7835
7840
7845
7850
7855
7860
7865
7870
7875
7880
7885
7890
7895
7900
7905
7910
7915
7920
7925
7930
7935
7940
7945
7950
7955
7960
7965
7970
7975
7980
7985
7990
7995
8000
8005
8010
8015
8020
8025
8030
8035
8040
8045
8050
8055
8060
8065
8070
8075
8080
8085
8090
8095
8100
8105
8110
8115
8120
8125
8130
8135
8140
8145
8150
8155
8160
8165
8170
8175
8180
8185
8190
8195
8200
8205
8210
8215
8220
8225
8230
8235
8240
8245
8250
8255
8260
8265
8270
8275
8280
8285
8290
8295
8300
8305
8310
8315
8320
8325
8330
8335
8340
8345
8350
8355
8360
8365
8370
8375
8380
8385
8390
8395
8400
8405
8410
8415
8420
8425
8430
8435
8440
8445
8450
8455
8460
8465
8470
8475
8480
8485
8490
8495
8500
8505
8510
8515
8520
8525
8530
8535
8540
8545
8550
8555
8560
8565
8570
8575
8580
8585
8590
8595
8600
8605
8610
8615
8620
8625
8630
8635
8640
8645
8650
8655
8660
8665
8670
8675
8680
8685
8690
8695
8700
8705
8710
8715
8720
8725
8730
8735
8740
8745
8750
8755
8760
8765
8770
8775
8780
8785
8790
8795
8800
8805
8810
8815
8820
8825
8830
8835
8840
8845
8850
8855
8860
8865
8870
8875
8880
8885
8890
8895
8900
8905
8910
8915
8920
8925
8930
8935
8940
8945
8950
8955
8960
8965
8970
8975
8980
8985
8990
8995
9000
9005
9010
9015
9020
9025
9030
9035
9040
9045
9050
9055
9060
9065
9070
9075
9080
9085
9090
9095
9100
9105
9110
9115
9120
9125
9130
9135
9140
9145
9150
9155
9160
9165
9170
9175
9180
9185
9190
9195
9200
9205
9210
9215
9220
9225
9230
9235
9240
9245
9250
9255
9260
9265
9270
9275
9280
9285
9290
9295
9300
9305
9310
9315
9320
9325
9330
9335
9340
9345
9350
9355
9360
9365
9370
9375
9380
9385
9390
9395
9400
9405
9410
9415
9420
9425
9430
9435
9440
9445
9450
9455
9460
9465
9470
9475
9480
9485
9490
9495
9500
9505
9510
9515
9520
9525
9530
9535
9540
9545
9550
9555
9560
9565
9570
9575
9580
9585
9590
9595
9600
9605
9610
9615
9620
9625
9630
9635
9640
9645
9650
9655
9660
9665
9670
9675
9680
9685
9690
9695
9700
9705
9710
9715
9720
9725
9730
9735
9740
9745
9750
9755
9760
9765
9770
9775
9780
9785
9790
9795
9800
9805
9810
9815
9820
9825
9830
9835
9840
9845
9850
9855
9860
9865
9870
9875
9880
9885
9890
9895
9900
9905
9910
9915
9920
9925
9930
9935
9940
9945
9950
9955
9960
9965
9970
9975
9980
9985
9990
9995
10000
10005
10010
10015
10020
10025
10030
10035
10040
10045
10050
10055
10060
10065
10070
10075
10080
10085
10090
10095
10100
10105
10110
10115
10120
10125
10130
10135
10140
10145
10150
10155
10160
10165
10170
10175
10180
10185
10190
10195
10200
10205
10210
10215
10220
10225

5 TG avoid tedious and inefficient procedures for DNA
isolation by directly proceeding to nucleic acid amplification
with minimal tissue handling. The greatest time saving measure is
the avoidance of DNA precipitation which in many protocols
10 requires overnight incubation and at least two repeat steps. Use
of tissue directly is based on the understanding that tissue DNA
is required only for the initial amplification cycles to generate
adequate amounts of DNA for subsequent amplification. The tissue
sample needs only to be optimally treated to render its DNA
15 available for initial amplification. By properly preparing the
topographic sample for direct use in the amplification reaction
an enormous savings in time and cost is realized.

20 Effective DNA amplification is the determining step for
successful genotyping. The creation of a storage unit which
results in a suspension of tissue particles of differing size and
character is essential to amplifiability. In addition
pretreatment steps as outlined below cause the topographic sample
to become further effectively amplified. For the majority of
cases pretreatment protocol 1 proves sufficient. In rare
25 instances inclusion of protocols 2 and 3 in that order meet the
need for effective amplification. Of importance in these protocols
is minimal sample manipulation for ease and timeliness of
application.

30 Conditions have been optimized and simplified for DNA
extraction while avoiding tedious nucleic acid precipitation. The
procedure is rapid and easy to perform with excellent
reproducibility and high efficiency for clinical practice and
research purposes. Specimens can sit overnight or up to a week at
room temperature in the Topographic lysis buffer (TLB) 1 without
any detriment allowing topographic sampling to be done at other
facilities and mailed (at room temperature) to a central

laboratory for further processing. Shorter incubations with higher concentrations of Proteinase K in TLB2 are equally effective, but overnight digestion has been found logistically to be the most convenient. Optimization of the Proteinase K

5 concentration with the incubation time is critical. If the Proteinase K is too concentrated or the incubation time is too long, the Proteinase K will digest the DNA that is of interest to be amplifying. Conversely a shorten incubation time or not as concentrated and the Proteinase K will not digest proteins
10 sufficient to render the DNA available. Tissue sampling tends to take place through the day and the sample filled tubes become available in the later afternoon for TLB mix adjustment. Following overnight digestion, the specimens are heated (99 ° Centigrade for 10 min.) and spun 2 min. in the morning at which
15 time they are then ready for the amplification process or freezer storage.

PROTOCOL 1: TOPOGRAPHIC SAMPLE PREPARATION, INITIAL TREATMENT

REAGENTS: Topographic Lysis Buffer 1 (100 mM NaCl, 25 mM EDTA pH 8.6, 10 mM Tris pH 8.3, 0.5% NP₄₀), and Topographic Lysis Buffer 2
20 (Proteinase K 2 mg/mL).

INSTRUMENTATION: microcentrifuge, pipettors and pipettes, water bath, -20° C freezer, 4° C refrigerator, tube racks, tube floats, thermal cycler.

Approximately 0.50 cm² of tissue requires 95 µl of Topographic
25 Lysis Buffer 1 and 5 µl of Topographic Lysis Buffer 2. To determine appropriate final volume of buffer mix, use following methods: (1) estimate tissue scrapped off slides; (2) make

comparable standards (scrape known quantities 0.50 cm², 0.25 cm² etc. into 25 µl TLB1.); or (3), experience. Samples are variable, but Topographic Genotyping system is very tolerant of small judgment errors. Specimens can sit up to a week in Topographic Lysis Buffer 1 without any detriment. Topographic Lysis Buffer 1 (TLB1) is 95% and Topographic Lysis Buffer 2 (TLB2) is 5% of total volume of Topographic Lysis Buffer Mix (TLBM). Remember 25 µl of Topographic Lysis Buffer 1 is already in sample tubes.

Add Topographic Lysis Buffer 1 and 2 to labeled tubes post-topographic selection (tissue in 25 µl TLB1). Example, 0.5 cm² requires additional 70 µl Topographic Lysis Buffer 1 and 5 µl Topographic Lysis Buffer 2, 0.25 cm² requires 22.5 µl TLB1 and 2.5 TLB2 for total volume of 50 µl. Additions are proportional.

Samples adjusted in TLB mix in step 1 are placed in a float in 42° C water bath for approximately 18 hr (overnight) for further digestion to extract DNA.

Tubes are removed and heated to 99° C for 10 min. (thermal cycler or water bath) to deactivate TLB2 Proteinase K and also to further disrupt cross linking of proteins.

Spin 2 min. at 12,000 rpm's (table top centrifuge) to pellet undigested proteins and membranes. Not every tube will have a pellet. Tubes are stored in -20° C freezer in labeled boxes in freezer racks and are ready for aliquoting into PCR reaction tube for DNA analysis.

DNA can be stored indefinitely. Specimen can be thawed, aliquots removed, and returned to freezer repeatedly for genetic analysis until depleted. Typically over 30 separate aliquots can be drawn

off for PCR reactions. After depletion, up to 20 μ l H₂O can be added and mixed in storage tube to wash DNA stuck on sides of tube into solution with H₂O. This sample can then be used successfully in more PCR reactions.

5 PROTOCOL 2: TOPOGRAPHIC SAMPLE PREPARATION, SECOND TREATMENT-
PHENOL CHLOROFORM TREATMENT

Unlike traditional Phenol/Chloroform treatment protocols, there is no ethanol precipitation step in which a DNA pellet is formed, dried (ethanol evaporated) and another step where fluid usually water is added to rehydrate the DNA pellet. Here, the sample remains in the same storage vial suspended in it's TLB mix and there is no change on the size of aliquot used in amplification protocol.

15 REAGENTS: Phenol:Chloroform:Isoamyl Alcohol (25:24:1),
Chloroform. (

20 INSTRUMENTATION: microcentrifuge, pipettors and pipettes, -20° C freezer, 4° refrigerator, tube racks.

Phenol Chloroform treatment starts at any step post overnight incubation in TLB mix, It can be done either before or after heating, centrifuging, or freezing storage steps. (Phenol deactivates Proteinase K). If this step is performed post overnight incubation, the heating step performed normally is not necessary because Phenol deactivated Proteinase K.

25 Equal volume of Phenol:Chloroform:Isoamyl Alcohol (25:24:1) is added to volume of sample in tube, mixed vigorously to a cream

color, then spun 30 sec. at 14,000 rpm's in centrifuge (room temp.) to separate aqueous phase (DNA in TLB mix) from Phenol phase (bottom) and interphase of proteins, membranes and other cell debris.

- 5 Open and draw off aqueous layer (DNA in TLB mix) into a new labeled tube discarding interface and phenol layer. Repeat step 1 and 2 again if necessary (rare).

Add equal volume of Chloroform, vortex 5 sec. and spin 30 sec. at 14,000 rpm's in centrifuge (room temp.). Draw off top aqueous layer from bottom Chloroform and add to new labeled tube. Repeat step 3 if there is any question of contamination of sample with Phenol (rare). Sample is ready for usual aliquoting to PCR reaction tube or freezer storage.

PROTOCOL 3: TOPOGRAPHIC TISSUE PREPARATION, THIRD TREATMENT-
REMOVAL OF SHORT LENGTH DNA FRAGMENTS

In a small minority of cases, topographic sample amplification proves ineffective following proteinase and phenol-chloroform treatment. Under these circumstance it is possible to effect nucleic acid amplification by filter separation of short length fragments of DNA. This effect is likely related to excessive nucleic acid breakdown in the same related to tissue breakdown prior to fixation, fixative effect, and or prolonged storage. These short DNA fragments may exert their negative effect on nucleic acid amplification by nonspecific triggering of the amplification reaction. Use of the filter is a simple step and does not involve tedious transferring of small liquid volumes.

The storage unit cellular suspension having been treated with proteinase K and phenol-chloroform is transferred into a Amicon centricon microtube filter system designed to produce a residual retentate of 25 microliters in which short DNA fragments less than 100 base pairs long has been permitted to pass through.

The sample is centrifuged until an appropriate retentate is produced. The sample may aliquoted for nucleic acid amplification or stored in this form.

NUCLEIC ACID AMPLIFICATION:

Nucleic acid amplification is carried out over 36 cycles in a 48 μ l volume. Sufficient product is generated to avoid need for internal nested primer amplification which carries a significant risk of contamination. Nucleic acid amplification is preferably carried out in a completely separate room away from the location where DNA sequencing is performed.

Optimal preliminary treatment of the tissue in the storage vial is critical to successful amplification of fixative treated DNA. Critical to tissue preparation is proper lysing of cell and nuclear membranes followed by appropriate protein digestion. Adjusting the PCR cycling parameters to favor poor quality DNA of fixative treated tissues is also critical for reliable results. Also valuable is careful addition of amplification enzyme at high temperature to minimize nonspecific amplification triggered by misplaced primer fragments. By careful handling of the storage tube it is possible to carry out numerous DNA sequence analysis in parallel from minute tissue targets.

PROTOCOL: NUCLEIC ACID AMPLIFICATION OF TOPOGRAPHIC SAMPLES

PRINCIPLE OF TEST: Area of selected gene is amplified from aliquot of storage sample by enzymatic nucleic acid amplification using primers (approx. 20 mer oligonucleotide) designed to hybridize to specific regions of the gene of interest.

REAGENTS: H₂O, MgCl₂, dNTP mix (A, G, C & T are mixed in equal volumes), primers, Taq polymerase, molecular biological grade mineral oil, Sample DNA from extraction procedure.

INSTRUMENTATION: Thermal cycler, dry bath (78° C), mini centrifuge, mixer, pipettors and tips, tube racks, permanent marker, tube opener, 4° C refrigerator, -20° Freezer.

Preparation of PCR mix and high temperature amplification onset format on the thermal cycler is done in a separate Pre-PCR lab to avoid contamination with amplified products in the Post-PCR lab. Use ultraviolet light when hood is not in use to break down DNA contamination. Amplification reactants mixing and aliquoting is done in hood. It is not necessary to work under a hood as long as work area is completely free from even aerosolized amplified products contaminants. After high temperature onset, the amplification reaction tubes are closed tight, cycled then removed to the Post amplification lab to be opened.

2. Remove buffer #2, dNTP's, and primers from -20° freezer to thaw. Remove H₂O and MgCl₂ from 4° C refrigerator. Mix gently and briefly centrifuge. Mineral oil is stored in hood. Turn on dry bath to heat.

3. Label 1.5 mL tubes for mixes. Each amplification reaction has its own Mix 1 tube labeled with its gene-exon. Mix 2 is unvarying (may be combined with one or more genes; when total volume is 1000 µl, split in two).

For formalin fixed tissue, cytology filter arcs and cytospin samples use following Mix 1: 18.5 µl H₂O

3.5 µl Buffer #2
7 µl MgCl₂
0.5 µl of each primer.

For cytology fluids, cell lines and fresh/frozen samples use the following variation of Mix 1: 23.5 μ l H₂O

3.5 μ l Buffer #2

2 μ l MgCl₂ 0.5

5 μ l of each primer.

All samples use Mix 2: 6.5 μ l H₂O

1.5 μ l Buffer #2

3 μ l MgCl₂

4 μ l dNTPs mix

(added later 0.25 μ l Taq Polymerase)

Place reagents on ice after their addition to the mixes and return to freezer as soon as possible.

4. Line storage samples in row of tube rack according to desired order. Add 30 μ l Mix 1 to each empty tube.

5. Add sample (filtered tips to avoid aerosol contamination) to corresponding tube. Formalin fixed tissue, cytology blocks and cytopsin samples in LB/PK are mixed and 3 μ l aliquoted out from bottom of tube (avoid pellet and floating debris). Cell lines and fresh/frozen samples in LB/PK are mixed and 1 μ l is aliquoted out from bottom of tube (avoid pellet). Filter arc is added and submerged. Cytology fluids are mixed and 5 μ l aliquoted into a pink tube to signal biohazard. (If sample depleted, up to 20 μ l of H₂O can be added, mixed in storage tube to wash DNA from sides and used successfully in PCR reaction.

6. Overlay reaction tubes with 2 drops (30 μ l) mineral oil, close and label with PCR run and sample # Label "primer only" tube with PCR # (negative control).

7. Tubes are placed in Thermal Cycler with times programmed as described below. Block will heat to 99° C for 5 min. count down. Taq is now removed from freezer and added to Mix 2, then mixed and centrifuged Return reagents to freezer and refrigerator. With 5 2 min. left in countdown, set Mix 2 in dry bath at 79° C.

8. After beep, Block will cool to 94° C for 5 min. count down. All tube lids are opened and 15 µl of Mix 2 are added to bottom, under mineral oil layer. Tubes are sealed and PCR lid shut. Cycling proceeds as programmed below.

9. After cycling has ended, block is at 4° C. Remove sealed tubes to sequencing lab for agarose electrophoresis. Store in 4° C for short time, or -20° C long term storage. Turn off PCR machine.

FILE A LINKED TO:

File: 99° C, 5 min., 1 cycle.

File: 94° C, 5 min., 1 cycle.

File: 98° C, 30 sec.; 55° C ,1 min.; 72° C, 3 min., 1 cycle.

File: 94° C, 1 min.; 55° C ,1 min.; 72° C, 5 min., 15 cycle.

File: 94° C, 1 min.; 55° C ,1 min.; 72° C, 10 min., 20 cycles.

File: 72° C, 15 min., 1 cycle.

File: 4 ° C soak.

25 To achieve specific and high yield gene amplification, delay addition of Taq polymerase until the entire reaction mixture is held at 94° C. This "hot start" is accomplished with addition of heated (79° C) Mix 2. Follow addition of Taq Polymerase with a

further high temperature ramp step of 30 seconds at 98° C for the first amplification cycle is important. This additional high temperature helps to overcome covalent bonding and limitation in motion related to fixation of tissue proteins and nucleic acids.

- 5 Allow longer times for individual steps. Avoid rapid ramping steps or shortened programs which have the undesirable effect of encouraging nonspecific product amplification at the expense of specific target gene amplification.

AMPLICON ISOLATION AND DNA SEQUENCING:

10 Agarose gel electrophoresis accomplishes two purposes. First it allows separation of oligonucleotide primers and other small sized nonspecific amplification product to be cleanly separated from amplicon. The isolation of pure true amplicon is essential for clean DNA sequencing. Second it enables evaluation of efficiency and specificity of amplification. The latter is 15 important in Quality Assurance and Control. Failed individual reactions may be repeated or undergo Phenol/Chloroform and filtering of short length DNA fragment treatment (see above).

20 Horizontal agarose electrophoresis is performed in standard fashion. TG produces sufficient desired product for a variety of DNA structural analyses including direct DNA sequencing. The latter, which results in detailed sequence information for comparative analysis is easily performed by standard protocols which isolate the amplified product from agarose following by 25 sequencing reactions such as dideoxy chain termination. Sufficient specific product is produced to enable repeated structural or sequence analyses from a single round of nucleic acid amplification.

Once amplified DNA has been obtained from the minute tissue sample, screening or specific detection of DNA can be performed to identify and characterize mutational damage. The most detailed analysis involves direct DNA sequencing which is usually performed using dideoxy chain termination and sulfur-35 radiolabelled nucleotide incorporation. (Sanger F, Nicklen S, Coulson AR, DNA Sequencing with Chain-Terminating Inhibitors. Proc. Natl. Acad. Sci. USA, 74:5463-5467, 1977), incorporated by reference. TG provides sufficient amplified DNA so that these procedures may be carried out under established standard conditions with high success. The most convenient format for clinical application would involve purification of amplified DNA from agarose followed by direct sequencing. The overall process from tissue receipt, histopathologic evaluation, target selection and topographic sampling, storage unit treatment, nucleic acid amplification and DNA sequence determination can be conveniently followed by the following computerized information system.

INFORMATICS

TG consists of a series of simple methodological steps beginning with tissue histology, proceeding through DNA amplification and mutational analysis and concluding with integration of morphologic and genetic information. The process of TG is broken down into parts in which several interrelated steps are performed by the technologist or pathologist/scientist. These parts usually consists of parallel manipulation of several individual specimens at the same time in order to achieve a significant savings in time and cost. Once a specimen has been accessioned and entered the TG analysis, the performance of individual technical components is performed blindly without knowledge concerning the biological significance of an individual sample. All specimens are therefore handled in an identical

fashion thereby avoiding introduction of bias. Only at the conclusion of the analysis, when integration of morphologic and genetic data is carried out, can the full significance of the individual sample made aware. To coordinate this effort an informatics system has been developed based upon the specific aspects of TG which functions to inform the user the progress and results of an single or groups of specimen samples.

This informatics program consists of a relational database which interrelates clinical, histopathologic, and genetic information (Table 3). TG informatics operates on eight levels of relationships consisting of over 75 fields which can be further configured to suit the individual needs of a particular user. This informatics database is very important in maintenance of quality control and assurance over the myriad activities of different parts of the analysis. The database also provides the means to determine trends and patterns in the integrated histopathologic and genetic information. For example the presence of certain genetic alterations in specific forms of human cancer can be made evident by analysis using the database and the statistical significance tested. TG informatic relational database is an essential component enabling the user to keep in touch with ongoing analysis, define genetic trends and maintain operating control.

As a further example, if a new cancer related genetic alteration is discovered, TG utilizes specific DNA sequence information to formulate appropriate nucleic acid amplifying primers and sequencing primers with which to identify and characterize the genetic mutation. TG allows the user to access any archival tissue specimens to best evaluate the significance of this potentially important genetic alteration. In and of itself this is of enormous advantage as the user is not

restricted to large quantity fresh or fresh-frozen tissue for testing. Instead the user could subject selected archival topographic specimens that have already been evaluated for genetic alterations. TG has the distinct attribute through the
5 creation of storage units to create a bank of readily available topographic samples circumventing the need to continuously recut tissue blocks for every new gene analysis. These storage units will already have been analyzed and therefore the new gene in question can be studied in the context of associated genetic
10 alterations on that specimen. In this way TG would have broad applicability greatly enhancing the information obtained to achieve assess in a more rapid fashion the significance of a newly identified genetic alteration.

The informatics programs outlined in Table 3 can be very
15 easily integrated with existing national, regional and local cancer registries. For example, patients with particular forms of cancer that have proven resistant to a newly introduced chemotherapy can be compared to patients with similar forms of cancer that exhibited a good response. The broad applicability of
20 TG would be ideal for such an analysis by its capacity to be effective on virtually any archival fixed tissue specimen irrespective of size or storage age.

The informatics program to support TG is outlined in Table
3. This application utilizes direct sequencing of the amplified
25 DNA to detect and characterize point mutational damage. In this format tissue histology, amplification, amplified product purification and DNA sequencing is coordinated by the informatics program as shown. The specific genetic targets currently being performed by TG is outlined in Table 3. This list is an under
30 representation of the applicability at this time as many more genetic targets could be incorporated in the analysis. TG

provides a means whereby new gene discoveries could be quickly brought to the clinical bedside and research laboratory.

4000270.110504

TABLE 3

INFORMATICS: OVERALL ORGANIZATION

	RELATIONAL LEVEL	FOCUS	NUMBER FIELDS
5	1	PATIENT	6
	2	TISSUE	10
	3	SPECIMEN	7
	4	STORAGE UNIT	9
	5	GENE	25
10	6	PCR AMPLIFICATION	8
	7	SEQUENCING GEL	9
	8	GENOTYPE	4

LEVEL 1: PATIENT RELATED DATA

	TITLE	ABBREVIATION	FIELD TYPE
15	LAST NAME	PT_LAST	TEXT
	FIRST NAME	PT_FIRST	TEXT
	MIDDLE INITIAL	PT_MID	TEXT
	SOCIAL SECURITY NO.	PT_SSN	NUMBER
	BIRTH DATE	PT_BIRTH	DATE
20	SEX	PT_SEX	TEXT
	COMMENT	PT_COM	TEXT

LEVEL 2: TISSUE RELATED DATA

	TITLE	ABBREV	FIELD TYPE
	SITE	TS_SITE1	TEXT
	ADDITIONAL SITE DESCRIPTION	TS_SITE2	TEXT
5	CELL	TS_CELL1	TEXT
	ADDITIONAL CELL DESCRIPTION	TS_CELL2	TEXT
	HISTOPATH	TS_HIST1	TEXT
	ADDITIONAL HISTOPATH DESCRIPTION	TS_HIST2	TEXT
10	PROGRESSION	TS_PROG1	TEXT
	ADDITIONAL PROGRESSION DESCRIPTION	TS_PROG2	TEXT
	PROJECT	TS_PROJ	TEXT
	COMMENTS	TS_COM	TEXT

15 LEVEL 3: SPECIMEN RELATED DATA

	TITLE	ABBREVIATION	FIELD TYPE
	DATE SPECIMEN TAKEN	SP_DATE	DATE
	PT AGE WHEN SPECIMEN TAKEN	SP_PTAGE	NUMBER
20	HOSPITAL	SP_HOSP	TEXT
	ACCESSION NUMBER	SP_ACCNO	TEXT
	BLOCK DESIGNATION	SP_BLOCK	TEXT
	TOPOGRAPHIC SITE	SP_TOP	TEXT
	FIXATION	SP_FIX	TEXT
25	COMMENT	SP_COM	TEXT

LEVEL 4: STORAGE UNIT

TITLE	ABBREV	FIELD TYPE
-------	--------	------------

	STORAGE LOCATION	ST_LOC	TEXT
	DATE TOPOGRAPHIC SELECTION	ST_DATE	DATE
	TOPOGRAPHER	ST_OPNA	TEXT
	NO. RECUTS USED	ST_NORC	NUMBER
5	NO. RECUTS UNDEPARAFFINED	ST_NOUND	NUMBER
	LEFT		
	NO RECUTS DEPARAFFINED LEFT	ST_NODEP	NUMBER
	NO. POST-TOPOGRAPHIC SLIDES	ST_NOPTT	NUMBER
	LEFT	G	
10	STORAGE UNIT TREATMENT	ST_TREAT	TEXT
	COMMENTS	ST_COM	TEXT

10000270110000

LEVEL 5: INDIVIDUAL GENE DATA

	TITLE	ABBREVIATIO N	FIELD TYPE
	H-RAS-1 EXON 1	GE_HRAS_01	TEXT
	H-RAS-1 EXON 2	GE_HRAS_02	TEXT
5	D2S123	GE_D2S123	TEXT
	A1AT EXON 2	GE_A1AT_02	TEXT
	A1AT EXON 3	GE_A1AT_03	TEXT
	A1AT EXON 5	GE_A1AT_05	TEXT
	GLUCOCEREBROSIDASE EXON 2	GE_GCS_02	TEXT
10	GLUCOCEREBROSIDASE EXON 9	GE_GCS_09	TEXT
	GLUCOCEREBROSIDASE EXON 10	GE_GCS_10	TEXT
	K-RAS-2 EXON 1	GE_KRAS_01	TEXT
	K-RAS-2 EXON 2	GE_KRAS_02	TEXT
	N-RAS EXON 1	GE_NRAS_01	TEXT
15	N-RAS EXON 2	GE_NRAS_02	TEXT
	P53 IMMUNOHISTOCHEMISTRY	GE_P53IMM	TEXT
	P53 EXON 5	GE_P53_01	TEXT
	P53 EXON 6	GE_P53_06	TEXT
	P53 EXON 7	GE_P53_07	TEXT
20	P53 EXON 8	GE_P53_08	TEXT
	MTS1 EXON 2 (PROX 1/3)	GE_MTS_02A	TEXT
	MTS1 EXON 2 (MID 1/3)	GE_MTS_02B	TEXT
	MTS1 EXON 2 (DIST 1/3)	GE_MTS_02C	TEXT
	C-RAF-1 EXON 15	GE_RAF_15	TEXT
25	C-RET EXON 7	GE_RET_07	TEXT
	C-RET EXON 8	GE_RET_08	TEXT
	HUMANPAPILLOMAVIRUS	GE_HPV	TEXT

LEVEL 6: NUCLEIC ACID AMPLIFICATION DATA

	TITLE	ABBREVIATION	FIELD TYPE
	PCR ACCESSION NUMBER	PCR_ACCNO	NUMBER
	PCR OPERATOR	PCR_OP	TEXT
5	PCR DATE PERFORMED	PCR_DATE	DATE
	PCR METHOD	PCR_METH	TEXT
	UPSTREAM PRIMER	PCR_UP	TEXT
	DOWNSTREAM PRIMER	PCR_DOWN	TEXT
	PCR RESULT	PCR_RSLT	TEXT
10	COMMENT	PCR_COM	TEXT

LEVEL 7: SEQUENCING GEL INFORMATION

	TITLE	ABBREVIATION	FIELD TYPE
	GEL ACCESSION NUMBER	GEL_ACCNO	NUMBER
	GEL OPERATOR	GEL_OP	NAME
15	GEL DATE PERFORMED	GEL_DATE	DATE
	SEQUENCING METHOD	GEL_SEQ	TEXT
	GEL METHOD	GEL_METH	TEXT
	SEQUENCING PRIMER	GEL_PRIM	TEXT
	RADIONUCLEOTIDE LABEL	GEL_LAB	TEXT
20	SEQUENCING RESULT	GEL_RSLT	TEXT
	COMMENT	GEL_COM	TEXT

LEVEL 8: GENOTYPING DATA

	TITLE	ABBREVIATION	FIELD TYPE
	SIGNIFICANCE	TG_SIG	TEXT
25	BASE RESULT	TG_BASE	TEXT
	AMINO ACID RESULT	TG_AA	TEXT
	ZYGOSITY & ALLELIC LOSS	TG_ZYG	TEXT

INFORMATICS DATABASE TEMPLATES

ARRANGEMENT OF FIELDS FOR COMPARATIVE DATA ANALYSIS

LOOKUP AN INDIVIDUAL PATIENT GENOTYPING RESULTS:

NAME|SSN|TISSUE PROJECT|HOSPITAL|ACCESSION NUMBER|BLOCK
5 DESIGNATION|TOPOGRAPHIC SITE|STORAGE LOCATION|PCR NO.|PCR
RESULT|GEL NO.|GEL RESULT|GENE|SIGNIFICANCE|BASE RESULT|AMINO
ACID RESULT|ZYGOSITY

PLANNING SPECIMEN SAMPLING:

TISSUE PROJECT|HOSPITAL|ACCESSION NUMBER|BLOCK
10 DESIGNATION|TOPOGRAPHIC SITE|DATE SPEC. TAKEN|PT AGE WHEN SPEC
TAKEN|FIXATION|SPECIMEN COMMENTS|NO. RECUTS UNDEPARAFFINIZED
LEFT

SETUP FOR TOPOGRAPHIC SELECTION:

15 STORAGE LOCATION|TISSUE PROJECT|HOSPITAL|ACCESSION NUMBER|BLOCK
DESIGNATION|TOPOGRAPHIC SITE|NO. RECUTS USED|NO. RECUTS
DEPARAFFINIZED LEFT|NO. POST-TOPOGRAPHIC SLIDES LEFT|STORAGE
COMMENTS

PCR SETUP:

20 GENE|TISSUE PROJECT|STORAGE LOCATION|PCR ACC. NO.|PCR
OPERATOR|PCR DATE|PCR METHOD|UPSTREAM PRIMER|DOWNSTREAM
PRIMER|PCR COMMENTS

PCR REVIEW:

25 PCR NO.|PCR RESULT|GENE|LAST NAME|SSN|ORGAN
SITE|HISTOPATHOLOGY|TISSUE PROJECT|HOSPITAL|ACCESSION
NUMBER|BLOCK DESIGNATION|TOPOGRAPHIC SITE|STORAGE LOCATION|GEL
ACC. NO.|GEL RESULT|BASE RESULT|AMINO ACID RESULT|ZYGOSITY|PCR
COMMENTS

SEQUENCING GEL SETUP:

GENE|TISSUE PROJECT|STORAGE LOCATION|GEL ACC. NO.|GEL
OPERATOR|GEL DATE|SEQUENCING METHOD|GEL METHOD|SEQUENCING
PRIMER|RADIONUCLEOTIDE LABEL|GEL COMMENTS

5 SEQUENCING GEL REVIEW:

GEL ACC. NO.|GEL RESULT|PCR NO.|GENE|LAST NAME|SSN|ORGAN
SITE|HISTOPATHOLOGY|TISSUE PROJECT|HOSPITAL|ACCESSION
NUMBER|BLOCK DESIGNATION|TOPOGRAPHIC SITE|STORAGE LOCATION|PCR
RESULT|BASE RESULT|AMINO ACID RESULT|ZYGOSITY|GEL COMMENTS

10 TISSUE/GENE ANALYSIS:

ORGAN SITE|ADD. ORGAN SITE|CELL TYPE|ADD. CELL
TYPE|HISTOPATH|ADD. HISTOPATH|PROGRESS|ADD. PROGRESS|TISSUE
PROJECT|GENE|SIGNIFICANCE|BASE RESULT|AMINO ACID
RESULT|ZYGOSITY

15 STATUS OF CASES IN PROGRESS:

PCR NO.|PCR RESULT|GEL NO.|GEL RESULT|TISSUE
PROJECT|GENE|SIGNIFICANCE|BASE RESULT|AMINO ACID
RESULT|ZYGOSITY

STATUS OF TISSUE PROJECT:

20 TISSUE PROJECT|NAME|HISTOPATH|HOSPITAL|ACC.
NO.|GENE|SIGNIFICANCE|BASE RESULT|AMINO ACID RESULT|ZYGOSITY

APPLICATION OF TOPOGRAPHIC GENOTYPING

TG is designed to enable fixative treated tissue specimens
to be sampled in minute detail according to microscopic cellular
25 characteristics so that the genetic analysis for specific DNA
damage is focused on the histopathologic properties of that
unique tissue sample. TG integrates genetic and histopathologic

analysis in a simple, timely, cost and includes a compatible relational database to track individual and groups of specimens and to evaluate aspects of quality control and assurance in a format suitably adapted for clinical application. With TG, the vast archive of fixative treated, paraffin-embedded tissue specimens becomes available for retrospective integrated genetic/histopathologic analysis. TG is equally effective irrespective of archival age or specimen size and as such may be expected to have very broad application. The use of TG is outlined in Table 4 and described below. The user of TG is free to investigate any tissue specimen for precise DNA sequence alterations.

MOLECULAR ONCOLOGY: Since the initial recognition of specific DNA coding mutational damage affecting certain oncogene and tumor suppresser genes, the list of such altered genes involved in human cancer has grown each year. Many currently studied genes consist of those shown Table 4 and may be expected to increase in number rapidly over time. TG provides the means whereby clinical tissue specimens may be critically evaluated for these important cancer related gene alterations.

The user of TG would decide upon a specific gene of interest to study for DNA structural alterations. He/she could consult any of a number of readily available repositories of genetic information such as GenBank, European Molecular Biology Library (EMBL), or other gene sequence database service. In addition to the sequence of the gene of interest such service provides information on gene structure including intron/exon splice sites, location of specific cancer related mutational damage and corresponding RNA and protein alterations. Alternatively, published manuscripts may be consulted for genetic sequence information of the oligonucleotide primers utilized to study a

specific gene target. Such information is all that is required to evaluate the presence, frequency, and significance of cancer related mutational injury. Amplification from fixative treated tissue is best limited by amplicon length to approximately 200
5 base pairs. Oligonucleotide primers would be fashioned based on genetic sequence information to center the amplification on the anticipated site of genetic damage. These oligonucleotide primers would be utilized for both DNA amplification and subsequent sequencing for detection and precise characterization of
10 mutational damage. The specificity of amplification would be confirmed by matching of provided DNA sequence information to that derived from the tissue specimen being evaluated.

Having appropriate oligonucleotide primers for the specific gene of interest TG would enable to the user to evaluate
15 informative tissue specimens for relationship to human cancer development and progression. At present the limitations incumbent upon the use of large fresh tissue specimens prevents analysis of optimal specimens for genetic study. TG would broaden the range of specimens available for evaluation thereby attaining critical
20 information at a much faster rate than that currently in practice. For example to determine whether a specific DNA sequence alteration was in fact predictive of degree of tumor aggressive or treatment responsiveness, recourse may be easily had to a tissue inventory in which specimens over 5 or 10 years
25 in age are recorded from patients who have been followed forward from that time. Such correlated tissue inventories are available from national, regional and local cancer registries whose function is to collate this type of follow-up information. The tissues in such registries is almost invariably fixative treated
30 and for many cancers would include minute sized specimens such as that obtained from biopsy sampling. Registries of this type are currently reluctant to provide specimens freely for molecular

analysis since current approaches result to total destruction of the tissue. TG would provide the means to use sparing amounts of such fixative treated tissues to yield critical genetic information. By more carefully selecting the tissue specimens from highly informative subjects which most critically evaluate a specific hypothesis, TG becomes the most effective means to attained causally related and predictive genetic information.

In addition to DNA coding mutational damage, TG can inform the user of the presence of gene allelic loss by examination of loss of heterozygosity. Loss of heterozygosity analysis (LOH) is an established genetic method in which the absence of one or both alleles from a tissue of interest is detected (Murphee AL, Benedict WF, Retinoblastoma: Clues to Human Oncogenesis. Science 223:1028-1033, 1985), incorporated by reference. Currently LOH is usually accomplished by restriction fragment length polymorphism analysis which requires relatively large sample size. TG, using nucleic acid amplification of microsatellite regions of DNA (Weissenbach J, A Second Generation Linkage Map of the Human Genome Based on Highly Informative Microsatellite Loci. Gene, 135:275-278, 1993), incorporated by reference, in proximity to genes lost during cancer progression, may be used to determine loss of heterozygosity. For loss of heterozygosity analysis by TG a sample of normal tissue and a separate sample to tumor tissue would be analyzed. The loss of one or both alleles from the tumor sample would indicate loss of genetic material in relationship to cancer growth and progression. At present, loss of heterozygosity determination is very often confounded by the fact that obtaining a large tissue specimen based on gross appearance only in the fresh state very often results in the undesired inclusion of normal tissue elements in significant amounts. TG avoids this complication by optimized purity of tissue elements in the sample based on careful histopathologic and topographic considerations

which control sample selection for genetic analysis. As described in the section on topographic sampling from glass slides, even when tumor is seen to infiltrate between normal cellular elements, by judicious wetting and scraping using the scalpel point it is possible to cleanly separate tumor from normal for selection of pure tumor sample. In this way it is possible not only to detect point mutational damage but to comment on the presence or absence of allelic loss.

TG for microsatellite analysis is one area of great potential as the number and location of microsatellites is very rapidly increasing due to the work of the Human Genome Project to fully map human DNA according to position of individual microsatellites. Genomic information sources such as GenBank may be electronically consulted for up to the minute detailed microsatellite maps and corresponding primer sequences upon which to base a loss of heterozygosity analysis for virtually any part of the human genome.

INHERITED GENETIC DISEASES: TG, by virtue of its capacity to effectively analyze archival fixed tissue, would be of enormous value in the diagnosis and characterization of inherited diseases. The list of genetic disease with known mutational change is enormous and expanding on a daily basis. Repositories of such information of major texts on the subject can be consulted through most of the network communication services such as Medline. Detection of these diseases is usually performed on a blood or other easily obtainable tissue sample such as skin. There are many instances however where this easy availability is not possible such as in the case when suspected family members are no longer alive. In these instances there may well be archival formalin fixed, paraffin embedded tissue specimens stored as an archival fixative treated tissue specimen in a

hospital laboratory. TG would represent the means to effectively analyze such tissue for informative genetic alterations.

TG would provide genetic information upon which to construct family tree type inheritance patterns in turn greatly increasing our general understanding of inherited genetic disease. Once a suspected genetic alteration was found in certain family members to be altered, recourse can be had to archival specimens to further evaluate the inheritance pattern of DNA sequence alterations prior to more broad based investigation of living family members. In this way a strong basis for the presence and type of genetic alterations can be achieved before living family members are involved in the evaluation. By searching archival tissue registries such as those available in most hospitals by specific clinical terminology, tissues from patients suspected for variants patterns of genetic disease may be identified and studied for their relationship to established disease states. TG expands the range of genetic analysis to include a broader cross section of patient tissue specimens better able to discover new and evaluate the role of specific genetic alterations.

GENETIC POLYMORPHISM DETERMINATION: In a similar fashion, TG can utilize microsatellite and genetic sequence polymorphism analysis as a tissue typing system for identification of individual subjects. Unique genetic markers would serve as the basis to clearly indicate that a specific tissue specimen in fact originates for a specific individual. This is of great value as a quality assurance measure in standard histopathology practice as it is not uncommon for small tissue specimens from one person may inadvertently contaminate the tissue specimens of another person. TG, being effective on fixative treated tissue specimens of small size, would represent the only means to specifically identify person of origin.

To accomplish this type of tissue identification, the user of TG would search genetic data banks for informative regions of the human genome for individual polymorphism. One such well established genetic targets are the HLA antigens, the genetic
5 sequence of which has been detailed for use in individual identification (Klein J, O'Huigin C, Composite Origin of Major Histocompatibility Complex Genes. Current Opinion in Genetics and Development, 3:923-930, 1993), incorporated by reference. The frequency with which such HLA targets show variation in the human
10 population have been described. Depending on the level of assurance desired, the user of TG would select one or more such targets for amplification. Identification can then be performed by direct sequencing or specific hybridization assay kits which are currently available. The feature of the storage unit in TG
15 would enable the user to obtain up to 20 to 30 or more individual assays upon which to build a detailed tissue identification analysis with great assurance. Furthermore this would all be accomplished without unnecessary sacrifice of the tissue block specimen which may be critically needed for histopathologic
20 evaluation.

TG could find similar use in forensic medicine when a need arises to relate a current tissue or blood specimen to that of an archival fixed tissue specimen. At present there is no effective means to consistently tissue type fixative treated material. TG
25 would provide then means to genotype and identify archival fixed tissue for comparison to other fresh or fixed tissue specimens.

INFECTIOUS DISEASES: TG is of enormous value in the diagnosis and characterization of microbial infection. Molecular methods currently are being applied to the detection of microbial
30 infection but in general requires a sizable fresh tissue specimen to be effective. TG would enable molecular detection to be

focused on the exact site or sites with a large complex tissue specimen wherein microorganisms may be expected to be found. Careful and precise selection of tissues for nucleic acid amplification would represent the key step to significantly
5 improving our ability to achieve earlier detection of microbial infection. Tissue selection for microorganism detection would follow microscopic evaluation, building upon the insights gained through histopathologic analysis, and not compete with morphologic study for precious tissue specimens.

10 For example, a tissue specimen may first be subject to careful histopathologic analysis wherein one or more minute abscess sites may be found. Such site would not be appreciated or
15 sampled when examining the gross specimen and therefore would be missed. Once found such minute abscess sites could be selectively sampled providing the enriched starting material for a nucleic acid/genotyping type of analysis for microorganism detection and
20 characterization. The approach offered by TG is particularly attractive since at present there is no other means to effectively deal with such a condition of microscopically discovered infection.

Even more importantly, TG would provide detailed structural information of the nature of a specific offending organism. Since amplification could be directed at any known region of the
25 organism information concerning mutational change or the presence of genomic virulence related factors would be available. TG would enable specific genotyping of microorganism based upon unique DNA content and therefore would represent an important ancillary with
30 which to detect and follow microbial infection. To accomplish this purpose the user may consult microbial genome sequence data banks such as those available through the National Institutes of Health. The information provided would allow the construction of

oligonucleotide probes designed to identify and characterized defined regions of the organism in question with particular emphasis on microbial virulence factors. By analyzing tissue specimens in a retrospective fashion from patients known to have pursued a certain course or to have exhibit a certain clinical pattern of treatment responsiveness, it becomes possible to use TG to establish the define the presence of new infectious agents or genotypes of microorganisms.

b6
b7C
b7D
b7E
b7F
b7G
b7H
b7I
b7J
b7K
b7L
b7M
b7N
b7O
b7P
b7Q
b7R
b7S
b7T
b7U
b7V
b7W
b7X
b7Y
b7Z
b7AA
b7AB
b7AC
b7AD
b7AE
b7AF
b7AG
b7AH
b7AI
b7AJ
b7AK
b7AL
b7AM
b7AN
b7AO
b7AP
b7AQ
b7AR
b7AS
b7AT
b7AU
b7AV
b7AW
b7AX
b7AY
b7AZ
b7BA
b7BB
b7BC
b7BD
b7BE
b7BF
b7BG
b7BH
b7BI
b7BJ
b7BK
b7BL
b7BM
b7BN
b7BO
b7BP
b7BQ
b7BR
b7BS
b7BT
b7BU
b7BV
b7BW
b7BX
b7BY
b7BZ
b7CA
b7CB
b7CC
b7CD
b7CE
b7CF
b7CG
b7CH
b7CI
b7CJ
b7CK
b7CL
b7CM
b7CN
b7CO
b7CP
b7CQ
b7CR
b7CS
b7CT
b7CU
b7CV
b7CW
b7CX
b7CY
b7CZ
b7DA
b7DB
b7DC
b7DD
b7DE
b7DF
b7DG
b7DH
b7DI
b7DJ
b7DK
b7DL
b7DM
b7DN
b7DO
b7DP
b7DQ
b7DR
b7DS
b7DT
b7DU
b7DV
b7DW
b7DX
b7DY
b7DZ
b7EA
b7EB
b7EC
b7ED
b7EE
b7EF
b7EG
b7EH
b7EI
b7EJ
b7EK
b7EL
b7EM
b7EN
b7EO
b7EP
b7EQ
b7ER
b7ES
b7ET
b7EU
b7EV
b7EW
b7EX
b7EY
b7EZ
b7FA
b7FB
b7FC
b7FD
b7FE
b7FF
b7FG
b7FH
b7FI
b7FJ
b7FK
b7FL
b7FM
b7FN
b7FO
b7FP
b7FQ
b7FR
b7FS
b7FT
b7FU
b7FV
b7FW
b7FX
b7FY
b7FZ
b7GA
b7GB
b7GC
b7GD
b7GE
b7GF
b7GG
b7GH
b7GI
b7GJ
b7GK
b7GL
b7GM
b7GN
b7GO
b7GP
b7GQ
b7GR
b7GS
b7GT
b7GU
b7GV
b7GW
b7GX
b7GY
b7GZ
b7HA
b7HB
b7HC
b7HD
b7HE
b7HF
b7HG
b7HH
b7HI
b7HJ
b7HK
b7HL
b7HM
b7HN
b7HO
b7HP
b7HQ
b7HR
b7HS
b7HT
b7HU
b7HV
b7HW
b7HX
b7HY
b7HZ
b7IA
b7IB
b7IC
b7ID
b7IE
b7IF
b7IG
b7IH
b7II
b7IJ
b7IK
b7IL
b7IM
b7IN
b7IO
b7IP
b7IQ
b7IR
b7IS
b7IT
b7IU
b7IV
b7IW
b7IX
b7IY
b7IZ
b7JA
b7JB
b7JC
b7JD
b7JE
b7JF
b7JG
b7JH
b7JI
b7JJ
b7JK
b7JL
b7JM
b7JN
b7JO
b7JP
b7JQ
b7JR
b7JS
b7JT
b7JU
b7JV
b7JW
b7JX
b7JY
b7JZ
b7KA
b7KB
b7KC
b7KD
b7KE
b7KF
b7KG
b7KH
b7KI
b7KJ
b7KK
b7KL
b7KM
b7KN
b7KO
b7KP
b7KQ
b7KR
b7KS
b7KT
b7KU
b7KV
b7KW
b7KX
b7KY
b7KZ
b7LA
b7LB
b7LC
b7LD
b7LE
b7LF
b7LG
b7LH
b7LI
b7LJ
b7LK
b7LL
b7LM
b7LN
b7LO
b7LP
b7LQ
b7LR
b7LS
b7LT
b7LU
b7LV
b7LW
b7LX
b7LY
b7LZ
b7MA
b7MB
b7MC
b7MD
b7ME
b7MF
b7MG
b7MH
b7MI
b7MJ
b7MK
b7ML
b7MM
b7MN
b7MO
b7MP
b7MQ
b7MR
b7MS
b7MT
b7MU
b7MV
b7MW
b7MX
b7MY
b7MZ
b7NA
b7NB
b7NC
b7ND
b7NE
b7NF
b7NG
b7NH
b7NI
b7NJ
b7NK
b7NL
b7NM
b7NN
b7NO
b7NP
b7NQ
b7NR
b7NS
b7NT
b7NU
b7NV
b7NW
b7NX
b7NY
b7NZ
b7OA
b7OB
b7OC
b7OD
b7OE
b7OF
b7OG
b7OH
b7OI
b7OJ
b7OK
b7OL
b7OM
b7ON
b7OO
b7OP
b7OQ
b7OR
b7OS
b7OT
b7OU
b7OV
b7OW
b7OX
b7OY
b7OZ
b7PA
b7PB
b7PC
b7PD
b7PE
b7PF
b7PG
b7PH
b7PI
b7PJ
b7PK
b7PL
b7PM
b7PN
b7PO
b7PP
b7PQ
b7PR
b7PS
b7PT
b7PU
b7PV
b7PW
b7PX
b7PY
b7PZ
b7QA
b7QB
b7QC
b7QD
b7QE
b7QF
b7QG
b7QH
b7QI
b7QJ
b7QK
b7QL
b7QM
b7QN
b7QO
b7QP
b7QQ
b7QR
b7QS
b7QT
b7QU
b7QV
b7QW
b7QX
b7QY
b7QZ
b7RA
b7RB
b7RC
b7RD
b7RE
b7RF
b7RG
b7RH
b7RI
b7RJ
b7RK
b7RL
b7RM
b7RN
b7RO
b7RP
b7RQ
b7RR
b7RS
b7RT
b7RU
b7RV
b7RW
b7RX
b7RY
b7RZ
b7SA
b7SB
b7SC
b7SD
b7SE
b7SF
b7SG
b7SH
b7SI
b7SJ
b7SK
b7SL
b7SM
b7SN
b7SO
b7SP
b7SQ
b7SR
b7SS
b7ST
b7SU
b7SV
b7SW
b7SX
b7SY
b7SZ
b7TA
b7TB
b7TC
b7TD
b7TE
b7TF
b7TG
b7TH
b7TI
b7TJ
b7TK
b7TL
b7TM
b7TN
b7TO
b7TP
b7TQ
b7TR
b7TS
b7TT
b7TU
b7TV
b7TW
b7TX
b7TY
b7TZ
b7UA
b7UB
b7UC
b7UD
b7UE
b7UF
b7UG
b7UH
b7UI
b7UJ
b7UK
b7UL
b7UM
b7UN
b7UO
b7UP
b7UQ
b7UR
b7US
b7UT
b7UU
b7UV
b7UW
b7UX
b7UY
b7UZ
b7VA
b7VB
b7VC
b7VD
b7VE
b7VF
b7VG
b7VH
b7VI
b7VJ
b7VK
b7VL
b7VM
b7VN
b7VO
b7VP
b7VQ
b7VR
b7VS
b7VT
b7VU
b7VV
b7VW
b7VX
b7VY
b7VZ
b7WA
b7WB
b7WC
b7WD
b7WE
b7WF
b7WG
b7WH
b7WI
b7WJ
b7WK
b7WL
b7WM
b7WN
b7WO
b7WP
b7WQ
b7WR
b7WS
b7WT
b7WU
b7WV
b7WW
b7WX
b7WY
b7WZ
b7XA
b7XB
b7XC
b7XD
b7XE
b7XF
b7XG
b7XH
b7XI
b7XJ
b7XK
b7XL
b7XM
b7XN
b7XO
b7XP
b7XQ
b7XR
b7XS
b7XT
b7XU
b7XV
b7XW
b7XX
b7XY
b7XZ
b7YA
b7YB
b7YC
b7YD
b7YE
b7YF
b7YG
b7YH
b7YI
b7YJ
b7YK
b7YL
b7YM
b7YN
b7YO
b7YP
b7YQ
b7YR
b7YS
b7YT
b7YU
b7YV
b7YW
b7YX
b7YY
b7YZ
b7ZA
b7ZB
b7ZC
b7ZD
b7ZE
b7ZF
b7ZG
b7ZH
b7ZI
b7ZJ
b7ZK
b7ZL
b7ZM
b7ZN
b7ZO
b7ZP
b7ZQ
b7ZR
b7ZS
b7ZT
b7ZU
b7ZV
b7ZW
b7ZX
b7ZY
b7ZZ

TABLE 4

APPLICATION OF TOPOGRAPHIC GENOTYPING

APPLICATION	EXAMPLE
MOLECULAR ONCOLOGY	
POINT MUTATIONAL DAMAGE	ONCOGENE/ANITONCOGENE MUTATIONAL CHANGE
ALLELIC LOSS	MICROSATELLITE LOSS OF HETEROZYGOSITY
MICROSATELLITE INSTABILITY	MICROSATELLITE MARKER INSTABILITY
INHERITED GENETIC DISEASE	GERMLINE MUTATIONAL CHANGE STUDY OF ARCHIVAL TISSUES FOR FAMILIAL INHERITANCE STUDIES
10 POLYMORPHISM DETERMINATION	GENOMIC POLYMORPHISM DETERMINATION MICROSATELLITE ANALYSIS QUALITY CONTROL OF CONTAMINATION IN PATHOLOGY

5

MICROORGANISM DETECTION	SPECIFIC MICROORGANISM DNA
MICROORGANISM	DETECTION
GENOTYPING	DNA BASED GENOTYPING
DETERMINATION OF	GENOMIC CORRELATES OF VIRULENCE
VIRULENCE	INTEGRATION RELATED DNA
VIRAL INTEGRATION	DETECTION

TABLE 5

CANCER RELATED DNA CODING GENE ALTERATIONS

GENE	GENE LOCATION	YEAR FIRS T DISC	AFFECTED HUMAN CANCERS
K-RAS-2	EXONS 1 & 2	1983	COLORECTAL, LUNG, PANCREAS ADENOCARCINOMA
H-RAS-1	EXONS 1 & 2	1983	GENITOURINARY & SKIN CANCERS
N-RAS	EXONS 1 & 2	1985	HEMATOLOGIC MALIGNANCIES
P53	EXONS 5-8	1988	MANY TYPES OF HUMAN CANCER
C-RET	EXONS 7 & 8	1988	THYROID & ENDOCRINE CANCERS
MCC	SCATTERED	1990	COLORECTAL & OTHER ADENOCARCINOMAS
APC	SCATTERED & MUTATION CLUSTER REG.	1990	COLORECTAL & OTHER ADENOCARCINOMAS

C-RAF-1	EXON 15	1993	LUNG CANCERS
VHL	EXONS 1-3	1993	RENAL CELL CARCINOMA VON HIPPEL LINDAU DISEASE
MTS1	EXONS 1 & 2	1994	MELANOMA & WIDE RANGE OF HUMAN CANCERS

Example of Use of Topographic Genotyping

5 The most frequent application of Topographic Genotyping (TG)
6 would be in the very common medical situation involving diagnosis
7 and treatment of human cancer. As described below, TG would be
8 used at several points throughout the patient course critically
9 influencing clinical decision-making. There is at present no
10 practical alternative means to obtain the information made
11 available by TG which can be had at minimal cost in a timely
12 fashion.

13 The patient with cancer would typically come to attention
14 when a mass lesion would be detected either directly as a
15 palpable nodule felt by the patient or indirectly by radiologic
16 examinations performed in response to specific clinical symptoms
17 or laboratory biochemical alterations. At this point, early in
18 the clinical course, the main objective is to establish a firm
19 diagnosis of the presence of cancer with minimal intervention
20 delaying major surgery until it can be planned and carried out
21 with greatest benefit. The usual practice is to obtain a small
22 sized biopsy specimen from the suspected site of cancer formation
23 and confirm the diagnosis by microscopic analysis. The small but
24 vital nature of the tissues specimens at this point in clinical
25 mamangement exclude most genetic analyses which consume

relatively large amounts of tissue preventing traditional histopathologic diagnosis.

TG is designed to work off minute targets sites from fixative treated tissue specimens providing specific information on the presence and type of cancer related gene damage present in that cancer. This is accomplished without any sacrifice a tissue or histologic treatment. In fact the molecular analysis provided by TG complements histopathologic evaluation confirming the presence of cancer and indicating the degree of biological aggressiveness inate that individual tumor. The integrated histopathologic/genetic information can be used in certain circumstances at this early stage to predict pattern and extent of cancer spread prior to it having even occurred. Patients can be placed into clinical subsets with greater or lesser risk for malignant progression. Through the use of TG on small sized biopsy speciemns the result is a more secure diagnosis of cancer and a better understanding of potential biological behavior on an individual case basis.

Once a decision to perform major surgery has been reached, TG would be intimately involved in the full laboratory analysis of the removed tissues. In the case of a mastectomy performed to excise a breast cancer, the first priority is to carefully and fully analyze the specimen for sites of cancer formation and spread and to characterize the malignancy according to traditional histopathologic features empirically known to predict tumor aggressiveness. Given the current emphasis on this disease and better methods for early diagnosis, many breast cancers are small in size. Despite the availability of a large tissue resection there still may be no fresh tissue remaining following histologic evaluation. More importantly detailed microscopic analysis is required in order to identify critical sites in the specimen

where genetic analysis needs to be applied. For example there may be two distinct sites of cancer formation as well as three separate sites where atypical cellular proliferations are found in the breast resection. These areas could only be identified by histologic examination and could not be sampled by grossly observing the specimen. TG is designed to use morphologic clues to sample the tissue at precise points of biological importance. As a result TG would be the essential means to analyze this type of specimen for detailed genetic alterations in turn used to guide medical management.

Later in the patient's clinical course, it may come to pass that tumor may recur at a distant site such as the liver. Having a history of breast cancer, the occurrence would be highly suspicious for return of breast cancer. This would nevertheless have to be confirmed which would be accomplished by once again obtaining a representative piece of tissue. Given the circumstances the tissue specimen would invariably be small in the form of a biopsy to histologically confirm cancer recurrence. Once again standard molecular technique would not be suitable due to the minute size of the sample. TG would provide the means to analyze the specimen and compare the genetic alterations to that present in the previous tissue specimens. This typical scenario points out how at three separate time TG would be the only effective means to analyze the tissue on a typical case of human cancer.

Finally the storage capability of TG permitting up to thirty separate genetic tests on a single topographic sample would allow newly discovered genetic alterations to be studied on previously sampled tissue specimens without the need to recut the tissue block. As our understanding of genetics and cancer increases together with a better knowledge of the human genome, TG would be

Although the invention has been described in detail in the foregoing embodiments for the purpose of illustration, it is to be understood that such detail is solely for that purpose and that variations can be made therein by those skilled in the art without departing from the spirit and scope of the invention except as it may be described by the following claims.

FULL-SCALE